



**A2 Mathematics Unit 3: Pure Mathematics B****Solutions and Mark Scheme**

| Question Number | Solution                                                                                                                                                                                             | Mark                                               | AO                                   | Notes                                                          |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------|--------------------------------------|----------------------------------------------------------------|
| 1. (a)          | $1 - \frac{x^2}{2} - 4x = x^2$ $\frac{3x^2}{2} + 4x - 1 = 0$ $3x^2 + 8x - 2 = 0$ $x = \frac{-8 \pm \sqrt{64 + 24}}{6} = \frac{-8 \pm \sqrt{88}}{6}$ $x = 0.230(1385...), (-2.896805...)$             | M1<br><br>A1<br><br>B1<br><br>B1<br><br><b>[4]</b> | AO1<br><br>AO1<br><br>AO1<br><br>AO1 | (Attempt to substitute for $\cos x, \sin x$ )<br><br>(Correct) |
| 2.              | $V = \frac{4}{3} \pi r^3$ $\frac{dV}{dt} = 3 \times \frac{4}{3} \pi r^2 \frac{dr}{dt}$ $4\pi \times 15^2 \frac{dr}{dt} = 250$ $\frac{dr}{dt} = \frac{250}{900\pi} \approx 0.088 \text{ (cm/second)}$ | <br><br>B1<br><br>M1<br><br>A1<br><br><b>[3]</b>   | <br><br>AO3<br><br>AO3<br><br>AO3    | <br><br><br><br>(Substitution of data)                         |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Mark                                   | AO                                          | Notes                                  |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------|---------------------------------------------|----------------------------------------|
| 3. (a)          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | G1<br>G1                               | AO1<br>AO1                                  | (Shape)<br>(Stationary point)          |
| (b) (i)         | A correct statement,<br>eg. $f^{-1}$ doesn't exist because $f$ is not a one-one function                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | E1                                     | AO2                                         |                                        |
| (ii)            | <p>Any appropriate domain<br/>eg. There are many possible appropriate domains. It is essential that any domain must be contained in one branch of the curve shown.</p> <p>Here we consider <math>(-3, \infty)</math>.</p> <p>Let <math>y = x^2 + 6x + 13</math><br/> <math>\quad = (x+3)^2 + 4</math></p> <p><math>x + 3 = \pm\sqrt{y-4}</math></p> <p>So that <math>x = -3 \pm \sqrt{y-4}</math></p> <p>Since <math>x &gt; -3</math>, the positive sign is appropriate</p> <p><math>\therefore x = -3 + \sqrt{y-4}</math></p> <p>And <math>f^{-1}(x) = -3 + \sqrt{x-4}</math></p> | B1<br><br>M1<br><br>A1<br>A1<br><br>A1 | AO2<br><br>AO1<br><br>AO1<br>AO2<br><br>AO2 | (Attempt to find $x$ in terms of $y$ ) |
|                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>[8]</b>                             |                                             |                                        |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Mark                                                            | AO                                                  | Notes                                                                |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------|-----------------------------------------------------|----------------------------------------------------------------------|
| 4. (a)          | $(1-x)^{-\frac{1}{2}} = 1 + \frac{x}{2} + \left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\frac{x^2}{2} + \dots$ $= 1 + \frac{x}{2} + \frac{3x^2}{8} + \dots$ <p>Valid for <math> x  &lt; 1</math></p><br><p>When <math>x = \frac{1}{10}</math>, <math>\left(\frac{9}{10}\right)^{-\frac{1}{2}} \approx 1 + \frac{1}{20} + \frac{3}{800} = \frac{843}{800}</math></p> <p>So that <math>(10)^{\frac{1}{2}} = 3 \times \frac{843}{800} = \frac{2529}{800}</math></p> | <p>B1</p> <p>B1</p><br><p>B1</p> <p>B1</p> <p><b>[4]</b></p>    | <p>AO1</p> <p>AO1</p><br><p>AO2</p> <p>AO1</p>      |                                                                      |
| 5.              | <p>After 30 years, saving is</p> $(1.08)1000 + (1.08)^2 1000 + \dots + (1.08)^{30} 1000$ <p>This is G.P with <math>a = (1.08)1000</math></p> $r = 1.08$ <p>and <math>n = 30</math></p> <p>Then</p> $S_{30} = (1000)(1.08) \left( \frac{(1.08)^{30} - 1}{0.08} \right)$ $\approx \text{£}122,346$                                                                                                                                                                     | <p>B1</p><br><p>B2</p><br><p>M1</p> <p>A1</p> <p><b>[5]</b></p> | <p>AO3</p><br><p>AO3, AO3</p> <p>AO3</p> <p>AO3</p> | <p>(B2 for 3 correct, B1 for 2 correct)</p> <p>(correct formula)</p> |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | Mark                                                                                             | AO                                                                                     | Notes                                    |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|------------------------------------------|
| 6.              | <p>If smallest side is <math>a</math>, largest side <math>= 8a</math></p> $8a = a + 14d$ $a = 2d$ $\text{Perimeter} = \frac{15}{2}[2a + 14d] = \frac{15}{2} \cdot 18d = 135d$ $\therefore 135d = 270$ $d = 2$ <p>Length of smallest side <math>= a = 2d = 4 \text{ cm}</math></p> <p><b>Alternative mark scheme:</b><br/> smallest side <math>= a</math>, largest side <math>= 8a</math></p> $\text{Perimeter} = \frac{15}{2}[a + 8a] = \frac{15}{2} \cdot 9a = \frac{135}{2}a$ $\therefore \frac{135}{2}a = 270$ $a = 4$ <p>Length of smallest side <math>= a = 4 \text{ cm}</math></p> | <p>M1<br/>A1</p> <p>M1</p> <p>B1</p> <p>(M1)<br/>(A1)</p> <p>(M1)<br/>(A1)</p> <p><b>[4]</b></p> | <p>AO3<br/>AO3</p> <p>AO3</p> <p>AO3</p> <p>(AO3)<br/>(AO3)</p> <p>(AO3)<br/>(AO3)</p> | <p>(Attempt to relate the two sides)</p> |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | Mark                                                       | AO                                                   | Notes                                                                                                                                                 |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| 7. (a)          | $\frac{d^2y}{dx^2} = 12ax^2 + 6bx + 36$ <p>For point of inflection at (1,11)</p> $12a + 6b + 36 = 0$ <p>So that <math>2a + b + 6 = 0</math> (1)</p>                                                                                                                                                                                                                                                                                                                                                               | M1                                                         | AO2                                                  | (attempt to find $\frac{d^2y}{dx^2}$ , 2 correct terms)                                                                                               |
| (b)             | <p>Also <math>a + b + 18 = 11</math> (2)</p> <p>From (1), (2), <math>a = 1</math>, <math>b = -8</math></p> $\therefore \frac{d^2y}{dx^2} = 12x^2 - 48x + 36$ $= 12(x^2 - 4x + 3) = 12(x-1)(x-3) = 0$ $\therefore \frac{d^2y}{dx^2} = 0 \text{ when } x = 3$ <p>and <math>\frac{d^2y}{dx^2}</math> changes sign as <math>x</math> passes through 3</p> <p><math>\therefore</math> There is a point of inflection at <math>x = 3</math>, <math>y = 3^4 - 8 \cdot 3^3 + 18 \cdot 3^2 = 27</math>, i.e at (3, 27)</p> | A1<br>B1<br>M1<br>A1<br><br>M1<br>A1<br><br>m1<br>A1<br>A1 | AO2<br>AO1<br>AO1<br>AO2<br>AO2<br>AO2<br>AO2<br>AO2 | (Attempt to solve for $a$ , $b$ )                                                                                                                     |
| (c)             | $\frac{dy}{dx} = 4x^3 - 24x^2 + 36x = 0$ $\therefore 4x(x^2 - 6x + 9) = 0$ <p>giving <math>x = 0, x = 3</math></p> <p>Then at <math>x = 0, y = 0</math> and <math>\frac{d^2y}{dx^2} = 36</math></p> <p>There is a minimum at <math>x = 0, y = 0</math></p>                                                                                                                                                                     | M2<br><br>A1<br><br>A1<br><br>G1<br>G1                     | AO1, AO1<br><br>AO1<br><br>AO1<br><br>AO1<br>AO1     | (M1 for correct differentiation but not equal to 0)<br>(point of Inflection)<br>(Two Values)<br><br>general shape<br><br>min two points of inflection |
|                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | [16]                                                       |                                                      |                                                                                                                                                       |

| Question Number | Solution                                                                                      | Mark        | AO       | Notes                              |
|-----------------|-----------------------------------------------------------------------------------------------|-------------|----------|------------------------------------|
| 8 (a) (i)       | $-\frac{e^{-3x+5}}{3} + C$                                                                    | M1          | AO1      | $(ke^{-3x+5})$                     |
|                 |                                                                                               | A1          | AO1      | $(k = -\frac{1}{3})$               |
|                 | (ii) $\int x^2 \ln x \, dx$                                                                   |             |          |                                    |
|                 | $u = \ln x, \frac{dv}{dx} = x^2$                                                              | M1          | AO1      | (Correct $u$ and $\frac{dv}{dx}$ ) |
|                 | $\frac{du}{dx} = \frac{1}{x}, v = \frac{x^3}{3}$                                              |             |          |                                    |
|                 | $\int x^2 \ln x \, dx = \frac{x^3}{3} \ln x - \int \frac{x^3}{3} \cdot \frac{1}{x} dx$        | A1,A1       | AO1, AO1 |                                    |
|                 | $= \frac{x^3}{3} \ln x - \frac{x^3}{9} + C$                                                   | A1          | AO1      |                                    |
|                 | (Penalise omission of C once only)                                                            |             |          |                                    |
|                 | (b) $\int_0^{\frac{1}{2}} \frac{x^2}{\sqrt{1-x^2}} dx$                                        |             |          |                                    |
|                 | $x = \sin \theta \quad dx = \cos \theta d\theta$                                              | B1          | AO3      |                                    |
|                 | $x = 0, \theta = 0 \quad x = \frac{1}{2}, \theta = \frac{\pi}{6}$                             | B1          | AO3      |                                    |
|                 | $= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta \cos \theta}{\sqrt{1-\sin^2 \theta}} d\theta$   | M1          | AO3      | (attempt to substitute)            |
|                 | $= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta$              | A1          | AO3      | (Correct)                          |
|                 | $= \int_0^{\frac{\pi}{6}} \sin^2 \theta d\theta$                                              | A1          | AO3      |                                    |
|                 | $= \int_0^{\frac{\pi}{6}} \frac{1 - \cos 2\theta}{2} d\theta$                                 | m1          | AO3      |                                    |
|                 | $= \left[ \frac{\theta}{2} - \frac{\sin 2\theta}{4} \right]_0^{\frac{\pi}{6}}$                | A1          | AO3      | (both correct)                     |
|                 | $\frac{\pi}{12} - \frac{\sin \frac{\pi}{3}}{4} - 0 + 0 = \frac{\pi}{12} - \frac{\sqrt{3}}{8}$ | A1          | AO3      |                                    |
|                 |                                                                                               | <b>[14]</b> |          |                                    |

| Question Number | Solution                                                           | Mark       | AO             | Notes                                            |
|-----------------|--------------------------------------------------------------------|------------|----------------|--------------------------------------------------|
| 9.              | $x^2 + 4 = 12 - x^2$                                               | M1         | AO3            | (Equating y's)                                   |
|                 | $2x^2 = 8$                                                         | A1         | AO3            |                                                  |
|                 | $x = \pm 2$                                                        |            |                |                                                  |
|                 | Area = $\int_{-2}^2 \{12 - x^2 - (x^2 + 4)\} dx$                   | M1         | AO3            | (expressing area)                                |
|                 | $= \int_{-2}^2 (8 - 2x^2) dx$                                      |            |                |                                                  |
|                 | $= \left[ 8x - \frac{2x^3}{3} \right]_{-2}^2$                      | A2         | AO3<br>AO3     | (F.T arithmetic error)                           |
|                 | $= \frac{64}{3}$                                                   | A1         | AO3            | (c.a.o)                                          |
|                 | <b>Alternative mark scheme for the Area:</b>                       |            |                |                                                  |
|                 | Area = $\int_{-2}^2 (12 - x^2) dx - \int_{-2}^2 (x^2 + 4) dx$      | (M1)       | (AO3)          |                                                  |
|                 | $= \left[ 12x - \frac{x^3}{3} - \frac{x^3}{3} - 4x \right]_{-2}^2$ | (A2)       | (AO3)<br>(AO3) | (A2 for 4 terms correct, A1 for 2 terms correct) |
|                 | $= \frac{64}{3}$                                                   | (A1)       | (AO3)          | (c.a.o)                                          |
|                 |                                                                    | <b>[6]</b> |                |                                                  |

| Question Number | Solution                                                                                                                                                                                                                                                                | Mark | AO  | Notes                                                                    |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|--------------------------------------------------------------------------|
| 10. (a)         | $f(x) = 1 + 5x - x^4$<br>$f(1) = 5, f(2) = -5$                                                                                                                                                                                                                          | M1   | AO2 | (Use of Intermediate Value Theorem.)<br>(correct values and conclusions) |
|                 | There is a change of sign indicating there is a root between 1 and 2.                                                                                                                                                                                                   | A1   | AO2 |                                                                          |
|                 | (b)<br>$x_{n+1} = \sqrt[4]{1 + 5x_n}, \quad x_0 = 1.5, \quad x_1 = 1.707476485$<br>$x_2 = 1.75734609$<br>$x_3 = 1.7687213, \quad x_4 = 1.7712854$<br>$x_5 = 1.771861948, \alpha \approx 1.77$                                                                           | B1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | B1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | B1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | B1   | AO1 |                                                                          |
|                 | (c)<br>$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{1 + 5x_n - x_n^4}{5 - 4x_n^3}$<br><br>$x_0 = 1.5$<br>$x_1 = 1.904411765$<br>$x_2 = 1.788115338$<br>$x_3 = 1.772305156$<br>$x_4 = 1.772029085$<br>$x_5 = 1.772028972$<br><br>Root $\alpha \approx 1.772029$ | M1   | AO1 | Attempt to use Newton-Raphson<br>All terms correct                       |
|                 |                                                                                                                                                                                                                                                                         | A1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | M1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | A1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         |      |     |                                                                          |
|                 |                                                                                                                                                                                                                                                                         |      |     |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | A1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | A1   | AO1 |                                                                          |
|                 |                                                                                                                                                                                                                                                                         |      |     | Correct to 6 decimal places                                              |
|                 |                                                                                                                                                                                                                                                                         |      |     |                                                                          |
|                 |                                                                                                                                                                                                                                                                         | [11] |     |                                                                          |

| Question Number | Solution                                                                         | Mark        | AO         | Notes                                          |
|-----------------|----------------------------------------------------------------------------------|-------------|------------|------------------------------------------------|
| 11. (a)         | $4x^3 + 2xy + x^2 \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$                          | B2          | AO1, AO1   | (B2, 4 correct terms)<br>(B1, 3 correct terms) |
|                 | Now, $x = -1, y = 3$                                                             |             |            |                                                |
|                 | so that $-4 - 6 + \frac{dy}{dx} + 6 \frac{dy}{dx} = 0$                           | B1          | AO1        |                                                |
|                 | $\frac{dy}{dx} = \frac{10}{7}$                                                   | B1          | AO1        |                                                |
|                 | (b) $\frac{dy}{dx} = \frac{dy}{dp} / \frac{dx}{dp} = \frac{2}{2p} = \frac{1}{p}$ | M1<br>A1    | AO1<br>AO1 |                                                |
|                 | Gradient of normal is $-p$                                                       | B1          | AO1        |                                                |
|                 | Equation of normal is $(y - 2p) = -p(x - p^2)$                                   | m1          | AO1        |                                                |
|                 | $y - 2p = -px + p^3$                                                             |             |            |                                                |
|                 | so that $y + px = 2p + p^3$                                                      | A1          | AO1        | convincing                                     |
|                 | When $y = 0, x = b$                                                              |             |            |                                                |
|                 | $b = 2 + p^2$                                                                    | B1          | AO2        |                                                |
|                 | Since $p^2 > 0, b > 2$                                                           | E1          | AO2        |                                                |
|                 |                                                                                  | <b>[11]</b> |            |                                                |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Mark                                              | AO                                                     | Notes                                         |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------|--------------------------------------------------------|-----------------------------------------------|
| 12. (a)         | <p>Let <math>y = \cos x</math></p> $\frac{dy}{dx} = \lim_{h \rightarrow 0} \left[ \frac{\cos(x+h) - \cos x}{h} \right]$ $= \lim_{h \rightarrow 0} \left[ \frac{\cos x \cos h - \sin x \sin h - \cos x}{h} \right]$ <p>As <math>h</math> approaches 0 <math>\cos h \approx 1 - \frac{h^2}{2}</math> and <math>\sin h \approx h</math></p> <p>So <math display="block">\frac{dy}{dx} = \lim_{h \rightarrow 0} \left[ \frac{\cos x \left( 1 - \frac{h^2}{2} \right) - \sin x \times h - \cos x}{h} \right]</math></p> $= \lim_{h \rightarrow 0} \left[ \frac{-\frac{h^2}{2} \cos x - h \sin x}{h} \right]$ $= -\sin x$ | <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> | <p>AO2</p> <p>AO2</p> <p>AO2</p> <p>AO2</p> <p>AO2</p> |                                               |
| (b) (i)         | $\frac{(x^3 + 1)6x - 3x^2(3x^2)}{(x^3 + 1)^2}$ $= \frac{3x(2 - x^3)}{(x^3 + 1)^2}$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | <p>M1</p> <p>A1</p>                               | <p>AO1</p> <p>AO1</p>                                  | (Correct formula)                             |
| (ii)            | $3x^2 \tan 3x + 3x^3 \sec^2 3x$ $= 3x^2 (\tan 3x + x \sec^2 3x)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <p>M1</p> <p>A1</p> <p><b>[9]</b></p>             | <p>AO1</p> <p>AO1</p>                                  | <p>(Correct formula)</p> <p>(All Correct)</p> |

| Question Number | Solution                                                                                                                                                                                                                                                                                            | Mark                                | AO                                          | Notes                                                                      |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|---------------------------------------------|----------------------------------------------------------------------------|
| 13. (a)         | $\operatorname{cosec}^2 x + \cot^2 x = 5$<br>$1 + 2 \cot^2 x = 5$<br>$\cot^2 x = 2$<br>$\tan x = \pm \frac{1}{\sqrt{2}}$<br>$x = 35.3, 215.3^\circ, 144.7^\circ, 324.7^\circ$                                                                                                                       | M1<br><br>A1<br><br>A1<br><br>B1,B1 | AO1<br><br>AO1<br><br>AO1<br><br>AO1<br>AO1 | (Attempt to write in terms of one function)<br><br><br><br><br>(each pair) |
| (b) (i)         | $4 \sin \theta + 3 \cos \theta \equiv R(\sin \theta \cos \alpha + \cos \theta \sin \alpha)$<br>$R \cos \alpha = 4$<br>$R \sin \alpha = 3$<br>$R = \sqrt{3^2 + 4^2} = 5$<br>$\tan \alpha = \frac{3}{4}, \alpha = 36.87^\circ$<br>$4 \sin \theta + 3 \cos \theta \equiv 5 \sin(\theta + 36.87^\circ)$ | B1<br>B1<br><br>B1<br>B1            | AO1<br>AO1<br><br>AO1<br>AO1                |                                                                            |
| (ii)            | $5 \sin(\theta + 36.87^\circ) = 2$<br>$\sin(\theta + 36.87^\circ) = 0.4$<br>$\theta + 36.87^\circ = 23.58^\circ, 156.42^\circ, 383.58^\circ$<br>$\theta = 119.5(5)^\circ, 346.7(1)^\circ$<br>$= 120^\circ, 347^\circ$ to the nearest degree                                                         | B1<br><br>B1<br><br>B1<br>B1        | AO1<br><br>AO1<br><br>AO1<br>AO1            |                                                                            |
|                 |                                                                                                                                                                                                                                                                                                     | <b>[12]</b>                         |                                             |                                                                            |

| Question Number | Solution                                                                                                                                                                                                                                                                                                                                                      | Mark                 | AO                       | Notes                                          |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------|--------------------------|------------------------------------------------|
| 14. (a)         | $\frac{dV}{dt} = 4 \frac{dh}{dt}$ $4 \frac{dh}{dt} = 0.004 - 0.0008 h$ $\frac{dh}{dt} = 0.001 - 0.0002 h$ $5000 \frac{dh}{dt} = 5 - h$                                                                                                                                                                                                                        | M1                   | AO3                      | (3 terms, at least 2 correct)                  |
| (b)             | $5000 \int \frac{dh}{5-h} = \int dt$ $-5000 \ln(5-h) = t + C \quad (1)$ $h = 0 \text{ at } t = 0$ $\therefore -5000 \ln(5) = C$ <p>Substitute in (1)</p> $-5000 \ln(5-h) = t - 5000 \ln(5)$ $t = 5000 \ln\left(\frac{5}{5-h}\right)$ $\therefore \left(\frac{5}{5-h}\right) = e^{\frac{t}{5000}}$ $5-h = 5e^{\frac{-t}{5000}}$ $h = 5 - 5e^{\frac{-t}{5000}}$ | M1<br>A1,A1<br>m1    | AO1<br>AO1<br>AO1        | (Separation of variables)<br>(-1 if C omitted) |
| (c)             | $h = 5 - 5e^{\frac{-3600}{5000}}$ $= 2.57 \text{ m}$                                                                                                                                                                                                                                                                                                          | A1<br>M1<br>A1<br>B1 | AO1<br>AO1<br>AO1<br>AO1 | (Attempt to invert)                            |
|                 |                                                                                                                                                                                                                                                                                                                                                               | <b>[10]</b>          |                          |                                                |

| Question Number | Solution                                                                                                                                                                            | Mark                                         | AO                            | Notes             |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|-------------------------------|-------------------|
| 15.             | $4x^2 + 9 < 12x$<br>$4x^2 - 12x + 9 < 0$<br>$(2x - 3)^2 < 0$<br>Impossible when $x$ is real.<br>Contradiction so that assumption is false.<br>$\therefore 4x + \frac{9}{x} \geq 12$ | M1<br><br><br>A1<br><br>A1<br><br><b>[3]</b> | AO2<br><br><br>AO2<br><br>AO2 | (Clear fractions) |