


## A2 Mathematics Unit 4: Applied Mathematics B

### Solutions and Mark Scheme

#### SECTION A – Statistics

Qu. No.	Solution	Mark	AO	Notes
1(a)	 <p>A = the event that a person has the disease. B = the event that a positive response is obtained</p> <p>Prob = <math>0.96 \times 0.99 = 0.9504</math></p> <p><b>Alternative mark scheme for (a):</b></p> <p>Prob = <math>0.96 \times 0.99</math> = 0.9504</p>	M1    A1  (M1) (A1)	AO1    AO2  (AO1) (AO2)	diagram
(b)	$P(B) = 0.04 \times 0.98 + 0.96 \times 0.01$ $= 0.0488$	M1 A1	AO3 AO1	
(c)	$P(A B) = \frac{P(A \cap B)}{P(B)}$ $= \frac{0.04 \times 0.98}{0.0488}$ $= 0.803(278688\dots)$	M1  A1  [6]	AO3  AO1	


Qu. No.	Solution	Mark	AO	Notes
2(a)(i)	$P(\text{J wins with 1}^{\text{st}} \text{ shot}) = P(\text{M misses}) \times P(\text{J hits})$ $= 0.75p$	M1 A1	AO1 AO1	
(ii)	<p>J wins with his second shot if the first three shots miss and then J hits the target with his second shot.</p> $P(\text{J wins with 2}^{\text{nd}} \text{ shot}) = 0.75 \times (1 - p) \times 0.75 \times p$	M1  A1	AO3  AO2	
(b)	$P(\text{J wins game}) = 0.75p + 0.75^2(1 - p)p + 0.75^3(1 - p)^2p + \dots$ <p>Attempting to sum an infinite geometric series</p> $= \frac{0.75p}{1 - 0.75(1 - p)}$ $= \frac{3p}{1 + 3p}$	M1  M1  A1	AO3  AO3  AO2	
(c)	<p>Mary is more likely to win if</p> $\frac{3p}{1 + 3p} < 0.5$ <p>leading to <math>p &lt; \frac{1}{3}</math></p>	M1  A1 <b>[9]</b>	AO3  AO1	
3(a)	<p>Continuous uniform distribution on [30,60]</p> <p>Mean = 45</p> <p>Variance = 75</p>	B1  B1 B1	AO3  AO1 AO1	
(b)	$P(\pi R^2 > 100) = P\left(R > \sqrt{\frac{100}{\pi}}\right)$ $= P\left(L > 2\pi\sqrt{\frac{100}{\pi}}\right)$ $= P(L > 35.45)$ $= \frac{60 - 35.45}{30} = 0.818(3) \text{ or } \frac{491}{600}$	M1  A1  A1 A1 <b>[7]</b>	AO3  AO2  AO1 AO1	

Qu. No.	Solution	Mark	AO	Notes
4(a)	Bell shaped	B1	AO2	Or Most values cluster in the middle of the range and the rest taper off symmetrically toward either extreme B0 for symmetrical only
(b)	$1 - P(6.12 < X < 8.12)$ $= 1 - 0.9949(0744)$ $= 0.0051$ (or 0.51%)	M1 A1	AO3 AO1	Or $P(X < 6.12) + P(X > 8.12)$ M1A0 For 0.9949(0744)
(c)(i)	The population of weights of 2p coins is normally distributed. Mean and median in the sample are very similar, suggesting a symmetric distribution.	B1 B1	AO2 AO2	B1B0 The weights of 2p coins are normally distributed. Population must be stated or implied.
(ii)	$H_0$ : The mean weight of all 2p coins in this batch = 7.12g $H_1$ : The mean weight of all 2p coins in this batch < 7.12g (one-sided)  $p\text{-value} = P(\bar{x} < 6.89 \mid H_0)$ $= P\left(z < \frac{6.89 - 7.12}{\frac{0.357}{\sqrt{30}}}\right)$ $= P(z < -3.52(874))$ $= 0.00021$ (allow 0.00022) Since $p\text{-value} < 0.01$ , Reject $H_0$  Very strong evidence to suggest the mean weight of the batch of 2p coins is less than 7.12(g)	B1  M1  A1 A1 A1  E1	AO3  AO1 AO1 AO2  AO3	Or $H_0: \mu = 7.12\text{g}$ B0 for $H_0$ : Mean = 7.12g Population must be stated or implied, ie. the batch of 2p coins  FT two-sided test $p\text{-value} = 2 \times 0.00021 = 0.00042$
	<b>Alternative Solution:</b>  $TS = \frac{6.89 - 7.12}{\frac{0.357}{\sqrt{30}}}$ $= -3.52(874)$ $CV = -2.32(63)$ Since $TS < CV$ Reject $H_0$  Very strong evidence to suggest the mean weight of the batch of 2p coins is less than 7.12(g)	(M1) (A1) (A1) (A1)	(AO1) (AO1) (AO1) (AO2)	FT Two-sided test $CVs = \pm 2.576$ Since $TS < -2.576$
		(E1)	(AO3)	
		[11]		

Qu. No.	Solution	Mark	AO	Notes
5(a)	$H_0: \rho = 0$ $H_1: \rho \neq 0$ two-sided	B1	AO3	$H_0: \rho = 0$ $H_1: \rho > 0$ one-sided Population stated or implied
	TS = 0.895	B1	AO1	TS = 0.895
	CV = $\pm 0.4821$	B1	AO1	CV = $\pm 0.412$
	Since TS > 0.4821, Reject $H_0$ Strong evidence to suggest the correlation coefficient is greater than zero	B1	AO2	Since TS > 0.412, Reject $H_0$
		E1	AO3	Strong evidence to suggest the correlation coefficient is greater than zero
(b)	P-value for correlation between Value for money and Cost per night is > 0.05	E1	AO2	
	Cost per night does not seem to be correlated to Value for money.	E1	AO2	
		<b>[7]</b>		

**SECTION B – Differential Equations and Mechanics**

Question Number	Solution	Mark	AO	Notes
6. (a)	$\mathbf{a} = \mathbf{F}/m = \frac{1}{4}(4\mathbf{i} - 12\mathbf{j})$ $\mathbf{a} = \mathbf{i} - 3\mathbf{j}$  Use $\mathbf{v} = \mathbf{u} + \mathbf{a}t$ , $\mathbf{u} = -\mathbf{i} + 4\mathbf{j}$ , $\mathbf{a} = \mathbf{i} - 3\mathbf{j}$ $\mathbf{v} = (-\mathbf{i} + 4\mathbf{j}) + 5(\mathbf{i} - 3\mathbf{j})$ $\mathbf{v} = 4\mathbf{i} - 11\mathbf{j}$	M1  M1 A1	AO3  AO2 AO1	
(b)	$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2 + 7\mathbf{i} - 26\mathbf{j}$  $\mathbf{s} = 2(-\mathbf{i} + 4\mathbf{j}) + \frac{1}{2} \times 4 \times (\mathbf{i} - 3\mathbf{j}) + (7\mathbf{i} - 26\mathbf{j})$ $\mathbf{s} = 7\mathbf{i} - 24\mathbf{j}$  $ \mathbf{s}  = \sqrt{7^2 + 24^2}$ $ \mathbf{s}  = 25$	M1  m1  A1  m1 A1 <b>[8]</b>	AO2  AO2  AO1  AO1 AO1	position vector relative to initial position vector. adding initial position vector.
7. (a)	Attempt to resolve in 2 directions  $T_1 \cos 23^\circ = T_2 \cos 40^\circ$ $T_1 \sin 23^\circ + T_2 \sin 40^\circ = 160$  Attempt to solve simultaneously  $T_1 = 137.56(028\dots)$ (N) $T_2 = 165.29(707\dots)$ (N)	M1  A1 A1  m1  A1 A1	AO3  AO2 AO2  AO1  AO1 AO1	dimensionally correct equation, no omitted or extra forces  correct equation correct equation  any valid method
(b)	Object modelled as particle Cable modelled as light strings	B1 B1  <b>[8]</b>	AO3 AO3	

Question Number	Solution	Mark	AO	Notes
8. (a)	$\frac{dP}{dt} = kP$ $\int \frac{dP}{P} = \int k dt$ $\ln P = kt + C$ <p>when <math>t = 0, P = 10</math></p> $C = \ln 10$ $\ln \frac{P}{10} = kt$ $e^{kt} = \frac{P}{10}$ $P = 10 e^{kt}$	M1  m1  A1  m1  m1  A1	AO3  AO2  AO1  AO2  AO1	separation of variables  correct integration
(b)	<p>When <math>t = 1, P = 20</math></p> $k = \ln 2$ $t = \frac{\ln 0.1P}{\ln 2}$ <p>When <math>P = 1000000</math></p> $t = \frac{\ln 1000000}{\ln 2}$ $t = 16.61 \text{ hours}$	M1    m1  A1 <b>[9]</b>	AO2    AO1  AO1	
9.	 <p> <math>R = mg = 12 \times 9.8 (= 117.6 \text{ N})</math>  Maximum friction <math>= \mu R</math>  Maximum friction <math>= 0.8 \times 12 \times 9.8</math>  <math>(= 94.08 \text{ N})</math> </p> <p>Therefore frictional force <math>= 75 \text{ (N)}</math>  because Max friction <math>&gt;</math> tractive force</p>	B1 M1 A1   B1 E1 <b>[5]</b>	AO1 AO3 AO1   AO3 AO3	used

Question Number	Solution	Mark	AO	Notes	
10.	(a)	$x = (V\cos\theta)t$	B1	AO1	
		$y = (V\sin\theta)t - \frac{1}{2}gt^2$	B1	AO1	
	(b)	$y = 0$ for time of flight	M1	AO2	
		$t = \frac{2V \sin \theta}{g}$			
		Range $R = V\cos\theta \cdot \frac{2V \sin \theta}{g}$	m1	AO2	
		$R = \frac{V^2 \sin 2\theta}{g}$	A1	AO2	
	(c) (i)	At maximum range, $\sin 2\theta = 1$	M1	AO3	oe
		$\theta = 45^\circ$			
		$\frac{V^2}{g} = 392$			
		$V = 62.0 \text{ (ms}^{-1}\text{)}$	A1	AO1	cao
	(ii)	$t = \frac{2 \times 62 \cdot 0 \times \sin 45}{g}$			
		$t = 8.95 \text{ (s)}$	A1	AO1	cao
	(iii)	Max height when $t = 4.47 \text{ s}$ ,	m1	AO2	
		$y_{max} = 62.5 \times \sin 45^\circ \times 4.47 - \frac{1}{2} \times 9.8 \times 4.47^2$			
		$y_{max} = 98.1 \text{ (m)}$	A1	AO1	cao
		[10]			