

C3

1.	0	0.5		
	0.2	0.401312339		
	0.4	0.310025518		
	0.6	0.231475216	(3 values correct)	B1
	0.8	0.167981614	(5 values correct)	B1

Correct formula with $h = 0.2$ M1

$$I \approx \frac{0.2}{3} \times \{0.5 + 0.167981614 + 4(0.401312339 + 0.231475216) + 2(0.310025518)\}$$

$$I \approx 0.2 \times 3.819182871 \div 3$$

$$I \approx 0.254612191$$

$$I \approx 0.2546 \quad (\text{f.t. one slip}) \quad \text{A1}$$

Note: Answer only with no working earns 0 marks

2.	(a)	e.g. $\theta = \frac{\pi}{2}$		
		$\cos \theta + \cos 4\theta = 1$	(choice of θ and one correct evaluation)	B1
		$\cos 2\theta + \cos 3\theta = -1$	(both evaluations correct but different)	B1

(b)	$2(\sec^2 \theta - 1) = \sec \theta + 8$	(correct use of $\tan^2 \theta = \sec^2 \theta - 1$)	M1
-----	--	---	----

An attempt to collect terms, form and solve quadratic equation in $\sec \theta$, either by using the quadratic formula or by getting the expression into the form $(a \sec \theta + b)(c \sec \theta + d)$,
with $a \times c = \text{coefficient of } \sec^2 \theta$ and $b \times d = \text{constant}$ m1

$$2 \sec^2 \theta - \sec \theta - 10 = 0 \Rightarrow (2 \sec \theta - 5)(\sec \theta + 2) = 0$$

$$\Rightarrow \sec \theta = \frac{5}{2}, \sec \theta = -2$$

$$\Rightarrow \cos \theta = \frac{2}{5}, \cos \theta = -\frac{1}{2} \quad (\text{c.a.o.}) \quad \text{A1}$$

$$\theta = 66.42^\circ, 293.58^\circ \quad \text{B1}$$

$$\theta = 120.0^\circ, 240.0^\circ \quad \text{B1 B1}$$

Note: Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.

$\cos \theta = +, -, \text{f.t.}$ for 3 marks, $\cos \theta = -, -, \text{f.t.}$ for 2 marks

$\cos \theta = +, +, \text{f.t.}$ for 1 mark

3.	(a)	$\frac{d(y^4)}{dx} = 4y^3 \frac{dy}{dx}$		B1
		$\frac{d(4x^2y)}{dx} = 4x^2 \frac{dy}{dx} + 8xy$		B1
		$\frac{d(3x^3 - 5x)}{dx} = 9x^2 - 5$		B1
		$\frac{dy}{dx} = \frac{9x^2 - 5 - 8xy}{4y^3 + 4x^2}$	(c.a.o.)	B1

(b)	$\frac{dx}{dt} = 4 - 2 \sin 2t,$	B1
	$\frac{dy}{dt} = 3 \cos 3t$	B1
	Use of $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$	M1
	Substituting $\frac{\pi}{12}$ for t in expression for $\frac{dy}{dx}$	m1
	$\frac{dy}{dx} = \frac{1}{\sqrt{2}}$	A1

4. $f(x) = 4x^3 - 2x - 5$
 An attempt to check values or signs of $f(x)$ at $x = 1, x = 2$ M1
 $f(1) = -3 < 0, f(2) = 23 > 0$
 Change of sign $\Rightarrow f(x) = 0$ has root in (1, 2) A1
 $x_0 = 1.2$
 $x_1 = 1.227601026$ (x_1 correct, at least 5 places after the point) B1
 $x_2 = 1.230645994$
 $x_3 = 1.230980996$
 $x_4 = 1.231017841 = 1.23102$ (x_4 correct to 5 decimal places) B1
 An attempt to check values or signs of $f(x)$ at $x = 1.231015, x = 1.231025$ M1
 $f(1.231015) = -1.197 \times 10^{-4} < 0, f(1.231025) = 4.218 \times 10^{-5} > 0$ A1
 Change of sign $\Rightarrow \alpha = 1.23102$ correct to five decimal places A1

Note: ‘Change of sign’ must appear at least once.

5. (a) (i) $\frac{dy}{dx} = 13 \times (7 + 2x)^{12} \times f(x), (f(x) \neq 1)$ M1
 $\frac{dy}{dx} = 26 \times (7 + 2x)^{12}$ A1
 (ii) $\frac{dy}{dx} = \frac{5}{\sqrt{1 - (5x)^2}}$ or $\frac{1}{\sqrt{1 - (5x)^2}}$ or $\frac{5}{\sqrt{1 - 5x^2}}$ M1
 $\frac{dy}{dx} = \frac{5}{\sqrt{1 - 25x^2}}$ A1
 (iii) $\frac{dy}{dx} = x^3 \times f(x) + e^{4x} \times g(x)$ M1
 $\frac{dy}{dx} = x^3 \times f(x) + e^{4x} \times g(x)$ (either $f(x) = 4e^{4x}$ or $g(x) = 3x^2$) A1
 $\frac{dy}{dx} = x^3 \times 4e^{4x} + e^{4x} \times 3x^2$ (all correct) A1

$$\begin{aligned}
 (b) \quad \frac{d}{dx}(\tan x) &= \frac{\cos x \times m \cos x - \sin x \times k \sin x}{\cos^2 x} \quad (m = \pm 1, k = \pm 1) & \text{M1} \\
 \frac{d}{dx}(\tan x) &= \frac{\cos x \times \cos x - \sin x \times (-\sin x)}{\cos^2 x} & \text{A1} \\
 \frac{d}{dx}(\tan x) &= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\
 \frac{d}{dx}(\tan x) &= \frac{1}{\cos^2 x} = \sec^2 x \quad (\text{convincing}) & \text{A1}
 \end{aligned}$$

$$\begin{aligned}
 6. \quad (a) \quad (i) \quad \int (7x-9)^{1/2} dx &= k \times \frac{(7x-9)^{3/2}}{3/2} + c \quad (k = 1, 7, 1/7) & \text{M1} \\
 \int (7x-9)^{1/2} dx &= 1/7 \times \frac{(7x-9)^{3/2}}{3/2} + c & \text{A1} \\
 (ii) \quad \int e^{x/6} dx &= k \times e^{x/6} + c \quad (k = 1, 6, 1/6) & \text{M1} \\
 \int e^{x/6} dx &= 6 \times e^{x/6} + c & \text{A1} \\
 (iii) \quad \int \frac{4}{5x-1} dx &= 4 \times k \times \ln |5x-1| + c \quad (k = 1, 5, 1/5) & \text{M1} \\
 \int \frac{4}{5x-1} dx &= 4 \times 1/5 \times \ln |5x-1| + c & \text{A1}
 \end{aligned}$$

$$\begin{aligned}
 (b) \quad \int (3x-4)^{-3} dx &= k \times \frac{(3x-4)^{-2}}{-2} \quad (k = 1, 3, 1/3) & \text{M1} \\
 \int_2^4 8 \times (3x-4)^{-3} dx &= \left[8 \times \frac{1}{3} \times \frac{(3x-4)^{-2}}{-2} \right]_2^4 & \text{A1}
 \end{aligned}$$

Correct method for substitution of limits M1


$$\int_2^4 8 \times (3x-4)^{-3} dx = \frac{5}{16} = 0.3125 \quad (\text{f.t. for } k = 1, 3 \text{ only}) \quad \text{A1}$$

7. (a) Trying to solve either $3x + 1 \leq 5$ or $3x + 1 \geq -5$ M1
 $3x + 1 \leq 5 \Rightarrow x \leq \frac{4}{3}$
 $3x + 1 \geq -5 \Rightarrow x \geq -2$ (both inequalities) A1
Required range: $-2 \leq x \leq \frac{4}{3}$ (f.t. one slip) A1

Alternative mark scheme

- $(3x + 1)^2 \leq 25$ (forming and trying to solve quadratic) M1
Critical points $x = -2$ and $x = \frac{4}{3}$ A1
Required range: $-2 \leq x \leq \frac{4}{3}$ (f.t. one slip in critical points) A1


- (b) (i)



Correct graph

B1

- (ii)



- Translation of graph of $f(x) = |x|$ with vertex at $(\pm 3, \pm 2)$ M1
Coordinates of vertex = $(3, 2)$ A1
Crosses y-axis at $(0, 5)$ A1

8. (a) $g'(x) = \frac{3 \times f(x)}{4x^2 + 9} + 2 \quad f(x) \neq 1$ M1
 $g'(x) = \frac{3 \times 8x}{4x^2 + 9} + 2$ A1
 $g'(x) = \frac{24x + 8x^2 + 18}{4x^2 + 9} = \frac{2(2x + 3)^2}{4x^2 + 9}$ (convincing) A1
- (b) (i) At stationary point, $\frac{2(2x + 3)^2}{4x^2 + 9} = 0$
or $\frac{3 \times 8x}{4x^2 + 9} + 2 = 0$ M1
 $\frac{2(2x + 3)^2}{4x^2 + 9} = 0$ only when $x = -\frac{3}{2}$ A1
- (ii) $g'(x) > 0$ either side of $x = -\frac{3}{2}$ (or at all other points) M1
Stationary point is a point of inflection A1
9. (a) $y - 5 = \ln(3x - 2)$ B1
An attempt to express candidate's equation as an exponential equation M1
 $x = \frac{(e^{y-5} + 2)}{3}$ (f.t. one slip) A1
 $f^{-1}(x) = \frac{(e^{x-5} + 2)}{3}$ (f.t. one slip) A1
- (b) $D(f^{-1}) = [5, \infty)$ B1
10. (a) $R(f) = [1, \infty)$ B1
 $R(g) = [-3, \infty)$ B1
- (b) $gf(x) = 2\sqrt{(x + 4)^2} - 3.$ M1
 $gf(x) = 2x + 5$ A1
- (c) $fg(x) = \sqrt{(2x^2 - 3 + 4)}$ (correct composition) B1
 $[fg(x)]^2 = 17^2$ (candidate's $fg(x)$) M1
 $x^2 = 144$ (f.t. one numerical slip) A1
 $x = \pm 12$ (c.a.o.) A1