



# **GCE MARKING SCHEME**

## **MATHEMATICS - C1-C4 & FP1-FP3 AS/Advanced**

**SUMMER 2013**

## INTRODUCTION

The marking schemes which follow were those used by WJEC for the Summer 2013 examination in GCE MATHEMATICS. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

<b>Paper</b>	<b>Page</b>
C1	1
C2	6
C3	11
C4	16
FP1	21
FP2	26
FP3	32

# C1

1. (a) (i) Gradient of  $BC = \frac{\text{increase in } y}{\text{increase in } x}$  M1  
 Gradient of  $BC = -4$  (or equivalent) A1  
 (ii) A correct method for finding the equation of  $BC$  using candidate's gradient for  $BC$  M1  
 Equation of  $BC$ :  $y - (-5) = -4(x - 6)$  (or equivalent) A1  
 (f.t. candidate's gradient of  $BC$ ) A1  
 Equation of  $BC$ :  $4x + y - 19 = 0$  (convincing) A1  
 (iii) Use of  $m_{AD} \times m_{BC} = -1$  M1  
 A correct method for finding the equation of  $AD$  using candidate's gradient for  $AD$  (M1)  
**(to be awarded only if corresponding M1 is not awarded in part (ii))**  
 Equation of  $AD$ :  $y - 4 = \frac{1}{4}(x - 8)$  (or equivalent) A1  
 (f.t. candidate's gradient of  $BC$ ) A1

**Note: Total mark for part (a) is 7 marks**

- (b) An attempt to solve equations of  $BC$  and  $AD$  simultaneously M1  
 $x = 4, y = 3$  (convincing) (c.a.o.) A1  
 (c) A correct method for finding the length of  $BD$  M1  
 $BD = \sqrt{68}$  A1  
 (d) A correct method for finding  $E$  M1  
 $E(0, 2)$  A1

2. (a)  $\frac{2 + 5\sqrt{7}}{4 + \sqrt{7}} = \frac{(2 + 5\sqrt{7})(4 - \sqrt{7})}{(4 + \sqrt{7})(4 - \sqrt{7})}$  M1  
 Numerator:  $8 - 2\sqrt{7} + 20\sqrt{7} - 35$  A1  
 Denominator:  $16 - 7$  A1  
 $\frac{2 + 5\sqrt{7}}{4 + \sqrt{7}} = -3 + 2\sqrt{7}$  (c.a.o.) A1

## Special case


If M1 not gained, allow B1 for correctly simplified numerator or denominator following multiplication of top and bottom by  $4 + \sqrt{7}$

- (b)  $\sqrt{360} = 6\sqrt{10}$  B1  
 $\sqrt{2} \times (\sqrt{5})^3 = 5\sqrt{10}$  B1  
 $\frac{\sqrt{30} \times \sqrt{8}}{\sqrt{6}} = 2\sqrt{10}$  B1  
 $\sqrt{360} - \sqrt{2} \times (\sqrt{5})^3 - \frac{\sqrt{30} \times \sqrt{8}}{\sqrt{6}} = -\sqrt{10}$  (c.a.o.) B1

3. (a)  $\frac{dy}{dx} = 4x - 10$  (an attempt to differentiate, at least one non-zero term correct) M1  
 An attempt to substitute  $x = 3$  in candidate's expression for  $\frac{dy}{dx}$  m1  
 Value of  $\frac{dy}{dx}$  at  $P = 2$  (c.a.o.) A1  
 Use of gradient of normal =  $\frac{-1}{\text{candidate's value for } \frac{dy}{dx}}$  m1  
 Equation of normal at  $P$ :  $y - (-5) = -\frac{1}{2}(x - 3)$  (or equivalent) A1  
 (f.t. candidate's value for  $\frac{dy}{dx}$  provided M1 and both m1's awarded) A1
- (b) An attempt to put candidate's expression for  $\frac{dy}{dx} = 0$  M1  
 $x$ -coordinate of  $Q = 2.5$   
 (f.t. one error in candidate's expression for  $\frac{dy}{dx}$ ) A1
4. (a)  $2(x - 4)^2 - 40$  B1 B1 B1
- (b) least value =  $-20$  (f.t. candidate's value for  $c$ ) B1  
 $x$ -coordinate =  $4$  (f.t. candidate's value for  $b$ ) B1
5. (a)  $(1 + 2x)^7 = 1 + 14x + 84x^2 \dots$  B1 B1 B1
- (b)  $(1 - 4x)(1 + 2x)^7 = 1 - 4x + 14x - 56x^2 + 84x^2$   
 Constant term and terms in  $x$  B1  
 Terms in  $x^2$  B1  
 (f.t. candidate's expression in (a))  
 $(1 - 4x)(1 + 2x)^7 = 1 + 10x + 28x^2$  (c.a.o.) B1

6. (a) (i) An expression for  $b^2 - 4ac$ , with at least two of  $a, b, c$  correct M1  
 $b^2 - 4ac = (4k + 1)^2 - 4 \times (k + 1) \times (k - 5)$  A1  
Putting  $b^2 - 4ac = 0$  m1  
 $4k^2 + 8k + 7 = 0$  (convincing) A1
- (ii) An expression for  $b^2 - 4ac$ , with at least two of  $a, b, c$  correct (M1)  
**(to be awarded only if corresponding M1 is not awarded in part (i))**  
 $b^2 - 4ac = 64 - 112 (= -48)$  A1  
 $b^2 - 4ac < 0 \Rightarrow$  no real roots A1
- Note: Total mark for part (a) is 6 marks**
- (b) Finding critical values  $x = -\frac{3}{4}, x = 3$  B1  
A statement (mathematical or otherwise) to the effect that  
 $x \leq -\frac{3}{4}$  or  $3 \leq x$  (or equivalent) B2  
(f.t. candidate's derived critical values)  
Deduct 1 mark for each of the following errors  
the use of strict inequalities  
the use of the word 'and' instead of the word 'or'
7. (a)  $y + \delta y = 5(x + \delta x)^2 + 8(x + \delta x) - 11$  B1  
Subtracting  $y$  from above to find  $\delta y$  M1  
 $\delta y = 10x\delta x + 5(\delta x)^2 + 8\delta x$  A1  
Dividing by  $\delta x$  and letting  $\delta x \rightarrow 0$  M1  
 $\frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} = 10x + 8$  (c.a.o.) A1
- (b)  $\frac{dy}{dx} = 6 \times \frac{2}{3} \times x^{-1/3} + 5 \times -2 \times x^{-3}$  (completely correct answer) B2  
**(If B2 not awarded, award B1 for at least one correct non-zero term)**
8. Attempting to find  $f(r) = 0$  for some value of  $r$  M1  
 $f(-1) = 0 \Rightarrow x + 1$  is a factor A1  
 $f(x) = (x + 1)(8x^2 + ax + b)$  with one of  $a, b$  correct M1  
 $f(x) = (x + 1)(8x^2 - 10x + 3)$  A1  
 $f(x) = (x + 1)(2x - 1)(4x - 3)$  (f.t. only  $8x^2 + 10x + 3$  in above line) A1  
 $x = -1, \frac{1}{2}, \frac{3}{4}$  (f.t. for factors  $2x \pm 1, 4x \pm 3$ ) A1

9. (a)



down curve with y-coordinate of maximum = 4  
 x-coordinate of maximum = -3  
 Both points of intersection with x-axis


B1

Concave

B1

B1

(b)



Concave down curve with y-coordinate of maximum = 4  
 x-coordinate of maximum = -1  
 Both points of intersection with x-axis

B1

B1

B1

Note: A candidate who draws a curve with no changes to the original graph is awarded 0 marks (both parts)

10. (a) (i)  $(2x \times x) + (2x \times x) + (2x \times y) + (2x \times y) + (x \times y) + (x \times y)$   
 $= 108$  M1  
 $6xy + 4x^2 = 108 \Rightarrow xy = 18 - \frac{2x^2}{3}$  (convincing) A1
- (ii)  $V = 2x \times x \times y = 2x(xy) \Rightarrow V = 36x - \frac{4x^3}{3}$  (convincing) B1
- (b)  $\frac{dV}{dx} = 36 - 3 \times \frac{4x^2}{3}$  B1  
 Putting derived  $\frac{dV}{dx} = 0$  M1  
 $x = 3, (-3)$  (f.t. candidate's  $\frac{dV}{dx}$ ) A1  
 Stationary value of  $V$  at  $x = 3$  is 72 (c.a.o) A1  
 A correct method for finding nature of the stationary point yielding a maximum value (for  $0 < x$ ) B1

## C2

1.	0	0.5	
	0.5	0.470588235	
	1	0.333333333	
	1.5	0.186046511	
	2	0.1	(5 values correct) B2
	(If B2 not awarded, award B1 for either 3 or 4 values correct)		

Correct formula with  $h = 0.5$  M1

$$I \approx \frac{0.5}{2} \times \{0.5 + 0.1 + 2(0.470588235 + 0.333333333 + 0.186046511)\}$$

$$I \approx 2.579936152 \times 0.5 \div 2$$

$$I \approx 0.644984038$$

$$I \approx 0.645 \quad (\text{f.t. one slip}) \quad \text{A1}$$

**Special case** for candidates who put  $h = 0.4$

0	0.5	
0.4	0.484496124	
0.8	0.398089172	
1.2	0.268240343	
1.6	0.164041994	
2	0.1	(all values correct) B1

Correct formula with  $h = 0.4$  M1

$$I \approx \frac{0.4}{2} \times \{0.5 + 0.1 + 2(0.484496124 + 0.398089172 + 0.268240343 + 0.164041994)\} \quad I \approx$$

$$3.229735266 \times 0.4 \div 2$$

$$I \approx 0.645947053$$

$$I \approx 0.646 \quad (\text{f.t. one slip}) \quad \text{A1}$$

**Note:** Answer only with no working earns 0 marks

2. (a) (i) Correct use of  $\tan \theta = \frac{\sin \theta}{\cos \theta}$  (o.e.) M1
- Correct use of  $\cos^2 \theta = 1 - \sin^2 \theta$  M1
- $6(1 - \sin^2 \theta) + 5 \sin \theta = 0 \Rightarrow 6 \sin^2 \theta - 5 \sin \theta - 6 = 0$  (convincing) A1
- (ii) An attempt to solve given quadratic equation in  $\sin \theta$ , either by using the quadratic formula or by getting the expression into the form  $(a \sin \theta + b)(c \sin \theta + d)$ ,  
with  $a \times c = 6$  and  $b \times d = -6$  M1
- $6 \sin^2 \theta - 5 \sin \theta - 6 = 0 \Rightarrow (3 \sin \theta + 2)(2 \sin \theta - 3) = 0$
- $\Rightarrow \sin \theta = -\frac{2}{3}, (\sin \theta = \frac{3}{2})$  (c.a.o.) A1
- $\theta = 221.81^\circ, 318.19^\circ$  B1 B1
- Note: Subtract (from final two marks) 1 mark for each additional root in range from  $3 \sin \theta + 2 = 0$ , ignore roots outside range.
- $\sin \theta = -$ , f.t. for 2 marks,  $\sin \theta = +$ , f.t. for 1 mark
- (b)  $2x - 60^\circ = -38^\circ, 38^\circ, 322^\circ$  (one value) B1
- $x = 11^\circ, 49^\circ$  B1 B1
- Note: Subtract (from final two marks) 1 mark for each additional root in range, ignore roots outside range.



3. (a) Either:  $(x+2)^2 = x^2 + (x-2)^2 - 2 \times x \times (x-2) \times \cos \hat{BAC}$   
 Or:  $\cos \hat{BAC} = \frac{x^2 + (x-2)^2 - (x+2)^2}{2 \times x \times (x-2)}$   
 (substituting the correct expressions in the correct places in the cos rule) M1  
 Either:  $\cos \hat{BAC} = \frac{x^2 + x^2 - 4x + 4 - x^2 - 4x - 4}{2 \times x \times (x-2)}$  (o.e.)  
 Or:  $\cos \hat{BAC} = \frac{x^2 + x^2 - 4x + 4 - x^2 - 4x - 4}{2x^2 - 4x}$  (o.e.) A1  
 $\cos \hat{BAC} = \frac{x-8}{2x-4}$  (convincing) A1
- (b) (i)  $\frac{x-8}{2x-4} = -\frac{1}{2}$  M1  
 $x = 5$  A1
- (ii) **Either:**  
 $\frac{\sin ABC}{3} = \frac{\sin 120^\circ}{7}$   
 (substituting the correct values in the correct places in the sin rule, f.t. candidate's value for  $x$ , provided  $x > 2$ ) M1  
 $ABC = 21.8^\circ$   
 (f.t. candidate's value for  $x$ , provided  $x > 2$ ) A1
- Or:**  
 $3^2 = 5^2 + 7^2 - 2 \times 5 \times 7 \times \cos ABC$   
 (substituting the correct values in the correct places in the cos rule, f.t. candidate's value for  $x$ , provided  $x > 2$ ) M1  
 $ABC = 21.8^\circ$   
 (f.t. candidate's value for  $x$ , provided  $x > 2$ ) A1
4. (a)  $S_n = a + [a + d] + \dots + [a + (n-1)d]$   
 (at least 3 terms, one at each end) B1  
 $S_n = [a + (n-1)d] + [a + (n-2)d] + \dots + a$   
 Either:  
 $2S_n = [a + a + (n-1)d] + [a + a + (n-1)d] + \dots + [a + a + (n-1)d]$   
 (at least three terms, including those derived from the first pair and the last pair plus one other pair of terms)  
 Or:  
 $2S_n = [a + a + (n-1)d] + \dots$  ( $n$  times) M1  
 $2S_n = n[2a + (n-1)d]$   
 $S_n = \frac{n}{2}[2a + (n-1)d]$  (convincing) A1

- (b) **Either:**
- $$\frac{10}{2}(2a + 9d) = 115 \quad \text{B1}$$
- $$S_{14} = 115 + 130 \quad \text{M1}$$
- $$\frac{14}{2}(2a + 13d) = 245 \quad \text{A1}$$
- An attempt to solve the candidate's equations simultaneously by eliminating one unknown M1
- $a = -2, d = 3$  (both values) (c.a.o.) A1
- Or:**
- $$\frac{10}{2}(2a + 9d) = 115 \quad \text{B1}$$
- $$(a + 10d) + (a + 11d) + (a + 12d) + (a + 13d) = 130 \quad \text{M1}$$
- $$4a + 46d = 130 \quad \text{(seen or implied by later work) A1}$$
- An attempt to solve the candidate's equations simultaneously by eliminating one unknown M1
- $a = -2, d = 3$  (both values) (c.a.o.) A1

5. (a)  $r = 0.8 \quad \text{B1}$
- $$S_{18} = \frac{100(1 - 0.8^{18})}{1 - 0.8} \quad \text{M1}$$
- $$S_{18} \approx 490.992 = 491 \quad \text{(c.a.o.) A1}$$
- (b) (i)  $ar = -20 \quad \text{B1}$
- $$\frac{a}{1 - r} = 64 \quad \text{B1}$$
- An attempt to solve these equations simultaneously by eliminating  $a$  M1
- $$16r^2 - 16r - 5 = 0 \quad \text{(convincing) A1}$$
- (ii)  $r = -\frac{1}{4} \quad \text{(c.a.o.) B1}$
- $$|r| < 1 \quad \text{E1}$$

6. (a)  $\frac{x^{5/4}}{5/4} + 2 \times \frac{x^{-4}}{-4} + c$  B1,B1  
(- 1 if no constant present)
- (b) (i)  $x^2 + 3 = 4x$  M1  
An attempt to rewrite and solve quadratic equation in  $x$ , either by using the quadratic formula or by getting the expression into the form  $(x + a)(x + b)$ , with  $a \times b = 3$  m1  
 $(x - 1)(x - 3) = 0 \Rightarrow x = 1, x = 3$  (both values, c.a.o) A1  
**Note: Answer only with no working earns 0 marks**
- (ii) Area of small triangle = 2  
(any method, f.t. candidate's value for  $x_A$ ) B1  
Use of integration to find the area under the curve M1  
 $\int x^2 dx = (1/3)x^3, \quad \int 3 dx = 3x$  (correct integration) B1  
Correct method of substitution of candidate's limits m1  
$$[(1/3)x^3 + 3x]_1^3 = (9 + 9) - (1/3 + 3) = 44/3$$
  
Use of candidate's values for  $x_A$  and  $x_B$  as limits and trying to find total area by adding area under curve to area of triangle m1  
Shaded area =  $44/3 + 2 = 50/3$  (c.a.o.) A1
7. (a) Let  $p = \log_a x, q = \log_a y$   
Then  $x = a^p, y = a^q$  (the relationship between log and power) B1  
 $xy = a^p \times a^q = a^{p+q}$  (the laws of indices) B1  
 $\log_a xy = p + q$  (the relationship between log and power)  
 $\log_a xy = p + q = \log_a x + \log_a y$  (convincing) B1
- (b) **Either:**  
 $(2 - 3x) \log_{10} 5 = \log_{10} 8$   
(taking logs on both sides and using the power law) M1  
 $x = \frac{2 \log_{10} 5 - \log_{10} 8}{3 \log_{10} 5}$  A1  
 $x = 0.236$  (f.t. one slip, see below) A1  
**Or:**  
 $2 - 3x = \log_5 8$  (rewriting as a log equation) M1  
 $x = \frac{2 - \log_5 8}{3}$  A1  
 $x = 0.236$  (f.t. one slip, see below) A1  
Note: an answer of  $x = -0.236$  from  $x = \frac{\log_{10} 8 - 2 \log_{10} 5}{3 \log_{10} 5}$   
earns M1 A0 A1  
an answer of  $x = 1.097$  from  $x = \frac{2 \log_{10} 5 + \log_{10} 8}{3 \log_{10} 5}$   
earns M1 A0 A1  
an answer of  $x = 0.708$  from  $x = \frac{2 \log_{10} 5 - \log_{10} 8}{\log_{10} 5}$   
earns M1 A0 A1

**Note: Answer only with no working shown earns 0 marks**

(c)	$\frac{1}{2} \log_a 144x^8 = \log_a 12x^4$	(power law)	B1
	$\log_a 90x^2 - \log_a \left[ \frac{5}{x} \right] = \log_a [90x^2 \times x]$	(subtraction law)	B1
	$\frac{90x^2 \times x}{5} = 12x^4$	(removing logs, f.t. one incorrect term)	B1
	$x = 1.5$	(c.a.o.)	B1
8.	(a)	$A(-1, 3)$	B1
		A correct method for finding the radius	M1
		Radius = 5	A1
	(b)	(i) Showing that the coordinates of A do not satisfy the equation of L (f.t. candidate's coordinates for A)	B1
		(ii) An attempt to substitute $(9 - x)$ for $y$ in the equation of $C_1$	M1
		$x^2 - 5x + 6 = 0$ (or $2x^2 - 10x + 12 = 0$ )	A1
		$x = 2, x = 3$	
		(correctly solving candidate's quadratic, both values)	A1
		Points of intersection are $(2, 7), (3, 6)$ (c.a.o.)	A1
	(c)	Distance between centres of $C_1$ and $C_2 = 13$	
9.		(f.t. candidate's coordinates for A)	B1
		Use of the fact that the shortest distance between the circles = distance between centres – sum of the radii	M1
		Shortest distance between the circles = 2	
		(f.t. candidate's coordinates for A and radius for $C_1$ .)	A1
	(a)	Substitution of values in area formula for triangle	M1
		Area = $\frac{1}{2} \times 7 \cdot 2^2 \times \sin 1.1 = 23.1 \text{ cm}^2$ .	A1
(b)		Let $\hat{BOC} = \phi$ radians	
		$\frac{1}{2} \times 7 \cdot 2^2 \times \phi = 19.44$	M1
		$\phi = 0.75$ (o.e.)	A1
		Length of arc $BC = 7.2 \times 0.75 = 5.4 \text{ cm}$	
			(f.t. candidate's value for $\phi$ ) A1

### C3

1. (a)
- |     |             |                    |    |
|-----|-------------|--------------------|----|
| 1   | 1.945910149 |                    |    |
| 1.5 | 2.238046572 |                    |    |
| 2   | 2.63905733  |                    |    |
| 2.5 | 3.073850053 |                    |    |
| 3   | 3.496507561 | (5 values correct) | B2 |
- (If B2 not awarded, award B1 for either 3 or 4 values correct)

Correct formula with  $h = 0.5$  M1  

$$I \approx \frac{0.5}{3} \times \{1.945910149 + 3.496507561 + 4(2.238046572 + 3.073850053) + 2(2.63905733)\}$$

$$I \approx 31.96811887 \times 0.5 \div 3$$

$$I \approx 5.328019812$$

$$I \approx 5.328 \quad \text{(f.t. one slip)} \quad \text{A1}$$

**Note:** Answer only with no working earns 0 marks

- (b)
- $$\int_1^3 \ln \sqrt{x^3 + 6} \, dx \approx 2.664 \quad \text{(f.t. candidate's answer to (a))} \quad \text{B1}$$

2. (a)
- $$4(\operatorname{cosec}^2 \theta - 1) - 8 = 2 \operatorname{cosec}^2 \theta - 5 \operatorname{cosec} \theta$$
- (correct use of  $\cot^2 \theta = \operatorname{cosec}^2 \theta - 1$ ) M1
- An attempt to collect terms, form and solve quadratic equation in  $\operatorname{cosec} \theta$ , either by using the quadratic formula or by getting the expression into the form  $(a \operatorname{cosec} \theta + b)(c \operatorname{cosec} \theta + d)$ , with  $a \times c = \text{coefficient of } \operatorname{cosec}^2 \theta$  and  $b \times d = \text{candidate's constant}$
- m1
- $$2 \operatorname{cosec}^2 \theta + 5 \operatorname{cosec} \theta - 12 = 0 \Rightarrow (2 \operatorname{cosec} \theta - 3)(\operatorname{cosec} \theta + 4) = 0$$
- $$\Rightarrow \operatorname{cosec} \theta = \frac{3}{2}, \operatorname{cosec} \theta = -4$$
- $$\Rightarrow \sin \theta = \frac{2}{3}, \sin \theta = -\frac{1}{4} \quad \text{(c.a.o.)} \quad \text{A1}$$
- $$\theta = 41.81^\circ, 138.19^\circ \quad \text{B1}$$
- $$\theta = 194.48^\circ, 345.52^\circ \quad \text{B1 B1}$$

Note: Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.

$\sin \theta = +, -, \text{ f.t. for 3 marks, } \sin \theta = -, -, \text{ f.t. for 2 marks}$   
 $\sin \theta = +, +, \text{ f.t. for 1 mark}$

- (b) Correct use of  $\sec \phi = \frac{1}{\cos \phi}$  and  $\tan \phi = \frac{\sin \phi}{\cos \phi}$  (o.e.) M1

$\sin \phi = -\frac{1}{2}$  A1  
 $\phi = 210^\circ, 330^\circ \quad \text{(f.t. for } \sin \phi = -a) \quad \text{A1}$

3. (a) Use of product formula yielding  $x^3 \times 2y \times \frac{dy}{dx} + 3x^2 \times y^2$  B1 B1  
 $\frac{dy}{dx} = -\frac{3x^2 y^2}{2x^3 y}$  (c.a.o.) B1
- (b) (i) Putting candidate's expression for  $\frac{dy}{dx} = 3$  and an attempt to simplify M1  
 $-\frac{3a^2 b^2}{2a^3 b} = 3 \Rightarrow b = -2a$  (convincing) A1  
(ii) Substituting  $a$  for  $x$  and  $-2a$  for  $y$  in the equation for  $C$  M1  
 $a = 2, b = -4$  A1
4. (a) Differentiating  $\ln t$  and  $5t^4$  with respect to  $t$ , at least one correct candidate's  $x$ -derivative  $= \frac{1}{t}$ , M1  
candidate's  $y$ -derivative  $= 20t^3$  (both values) A1  
 $\frac{dy}{dx} = \frac{\text{candidate's } y\text{-derivative}}{\text{candidate's } x\text{-derivative}}$  M1  
 $\frac{dy}{dx} = 20t^4$  (c.a.o.) A1
- (b)  $\frac{d}{dt} \left[ \frac{dy}{dx} \right] = 80t^3$  (f.t. candidate's expression for  $\frac{dy}{dx}$ ) B1  
Use of  $\frac{d^2 y}{dx^2} = \frac{d}{dt} \left[ \frac{dy}{dx} \right] \div \text{candidate's } x\text{-derivative}$  M1  
 $\frac{d^2 y}{dx^2} = 80t^4$  (f.t. one slip) A1  
 $\frac{d^2 y}{dx^2} = 0.648 \Rightarrow t = 0.3$  (c.a.o.) A1
5. (a)  $\frac{dy}{dx} = 5 \times (7 - 9x^2)^4 \times f(x),$  ( $f(x) \neq 1$ ) M1  
 $\frac{dy}{dx} = -90x \times (7 - 9x^2)^4$  A1
- (b)  $\frac{dy}{dx} = \frac{6}{1 + (6x)^2}$  or  $\frac{1}{1 + (6x)^2}$  or  $\frac{6}{1 + 36x^2}$  M1  
 $\frac{dy}{dx} = \frac{6}{1 + 36x^2}$  A1
- (c)  $\frac{dy}{dx} = e^{4x} \times m \sec^2 2x + \tan 2x \times k e^{4x}$  ( $m = 1, 2, k = 1, 4$ ) M1  
 $\frac{dy}{dx} = e^{4x} \times 2 \sec^2 2x + \tan 2x \times 4e^{4x}$  (at least one correct term) B1  
 $\frac{dy}{dx} = e^{4x} \times 2 \sec^2 2x + \tan 2x \times 4e^{4x}$  (c.a.o.) A1

$$(d) \quad \frac{dy}{dx} = \frac{(2 + \cos x) \times m \cos x - (3 + \sin x) \times k \sin x}{(2 + \cos x)^2} \quad (m = 1, -1 \quad k = 1, -1) \quad \text{M1}$$

$$\frac{dy}{dx} = \frac{(2 + \cos x) \times (\cos x) - (3 + \sin x) \times (-\sin x)}{(2 + \cos x)^2} \quad \text{A1}$$

$$\frac{dy}{dx} = \frac{2 \cos x + 3 \sin x + 1}{(2 + \cos x)^2} \quad \text{A1}$$

$$6. \quad (a) \quad (i) \quad \int \cos(3x + \pi/2) dx = k \times \sin(3x + \pi/2) + c \quad (k = 1, 3, 1/3, -1/3) \quad \text{M1}$$

$$\int \cos(3x + \pi/2) dx = 1/3 \times \sin(3x + \pi/2) + c \quad \text{A1}$$

$$(ii) \quad \int e^{3-4x} dx = k \times e^{3-4x} + c \quad (k = 1, -4, 1/4, -1/4) \quad \text{M1}$$

$$\int e^{3-4x} dx = -1/4 \times e^{3-4x} + c \quad \text{A1}$$

$$(iii) \quad \int \frac{7}{8x+5} dx = 7 \times k \times \ln|8x+5| + c \quad (k = 1, 8, 1/8) \quad \text{M1}$$

$$\int \frac{7}{8x+5} dx = 7 \times 1/8 \times \ln|8x+5| + c \quad \text{A1}$$

**Note: The omission of the constant of integration is only penalised once.**

$$(b) \quad \int (2x-1)^{-4} dx = k \times \frac{(2x-1)^{-3}}{-3} \quad (k = 1, 2, 1/2) \quad \text{M1}$$

$$\int_1^2 9 \times (2x-1)^{-4} dx = \left[ 9 \times 1/2 \times \frac{(2x-1)^{-3}}{-3} \right]_1^2 \quad \text{A1}$$

Correct method for substitution of limits in an expression of the form  $m \times (2x-1)^{-3}$   
M1

$$\int_1^2 9 \times (2x-1)^{-4} dx = \frac{13}{9} = 1.44 \quad (\text{f.t. for } k = 1, 2 \text{ only}) \quad \text{A1}$$

**Note: Answer only with no working earns 0 marks**


7. (a) Choice of  $a \neq -1$  and  $b = -a - 2$  M1  
 Correct verification that given equation is satisfied A1
- (b) Trying to solve either  $x^2 - 10 \leq 6$  or  $x^2 - 10 \geq -6$  M1  
 $x^2 - 10 \leq 6 \Rightarrow x^2 \leq 16$   
 $x^2 - 10 \geq -6 \Rightarrow x^2 \geq 4$  (both inequalities) A1  
 At least one of:  $2 \leq x \leq 4, -4 \leq x \leq -2$  (f.t. one slip) A1  
 Required range:  $2 \leq x \leq 4$  or  $-4 \leq x \leq -2$  (c.a.o.) A1

**Alternative mark scheme**

- $(x^2 - 10)^2 \leq 36$  (forming and trying to solve quadratic in  $x^2$ ) M1  
 Critical values  $x^2 = 4$  and  $x^2 = 16$  A1  
 At least one of:  $2 \leq x \leq 4, -4 \leq x \leq -2$  (f.t. one slip) A1  
 Required range:  $2 \leq x \leq 4$  or  $-4 \leq x \leq -2$  (c.a.o.) A1

8.  $x_0 = -1.5$   
 $x_1 = -1.666394263$  ( $x_1$  correct, at least 5 places after the point) B1  
 $x_2 = -1.676625462$   
 $x_3 = -1.677198866$   
 $x_4 = -1.677230823 = -1.67723$  ( $x_4$  correct to 5 decimal places) B1  
 Let  $f(x) = x^2 + e^x - 3$   
 An attempt to check values or signs of  $f(x)$  at  $x = -1.677225, x = -1.677235$  M1  
 $f(-1.677225) = -2.44 \times 10^{-5} < 0, f(-1.677235) = 7.26 \times 10^{-6} > 0$  A1  
 Change of sign  $\Rightarrow \alpha = -1.67723$  correct to five decimal places A1

9.



- Concave down curve and y-coordinate of maximum = 4 B1  
 x-coordinate of maximum = -1 B1  
 Both points of intersection with x-axis B1



10. (a)  $y - 6 = e^{5 - x/2}$ . B1  
 An attempt to express equation as a logarithmic equation and to isolate  $x$  M1  
 $x = 2 [5 - \ln (y - 6)]$  (c.a.o.) A1  
 $f^{-1}(x) = 2 [5 - \ln (x - 6)]$   
 (f.t. one slip in candidate's expression for  $x$ ) A1
- (b)  $D(f^{-1}) = [7, \infty)$  B1 B1
11. (a) (i)  $D(fg) = (0, \pi/4]$  B1  
 (ii)  $R(fg) = (-\infty, 0]$  B1 B1
- (b) (i)  $fg(x) = -0.4 \Rightarrow \tan x = e^{-0.4}$  M1  
 $x = 0.59$  A1  
 (ii) Equation has solution only if  $k \in R(fg)$ .  
 $\therefore$  choose any  $k \notin R(fg)$  (f.t. candidate's  $R(fg)$ ) B1

# C4

1. (a)  $f(x) \equiv \frac{A}{x^2} + \frac{B}{x} + \frac{C}{(x+2)}$  (correct form) M1

$$6 + x - 9x^2 \equiv A(x+2) + Bx(x+2) + Cx^2$$

(correct clearing of fractions and genuine attempt to find coefficients)

m1

$$A = 3, C = -8, B = -1 \quad (\text{all three coefficients correct}) \quad \text{A2}$$

If A2 not awarded, award A1 for at least one correct coefficient

(b) (i)  $f'(x) = \frac{-6}{x^3} + \frac{1}{x^2} + \frac{8}{(x+2)^2}$  (o.e.)

(f.t. candidate's values for A, B, C)

(first term)

B1

(at least one of last two terms)

B1

(ii)  $f'(2) = 0 \Rightarrow$  stationary value when  $x = 2$  (c.a.o.) B1

2.  $3x^2 - 2x \times 2y \frac{dy}{dx} - 2y^2 + 3y^2 \frac{dy}{dx} = 0$   $\left[ \begin{array}{l} -2x \times 2y \frac{dy}{dx} - 2y^2 \\ \frac{dy}{dx} \end{array} \right]$  B1

$$\left[ \begin{array}{l} 3x^2, 3y^2 \frac{dy}{dx} \\ \frac{dy}{dx} \end{array} \right]$$
 B1

**Either**  $\frac{dy}{dx} = \frac{2y^2 - 3x^2}{3y^2 - 4xy}$  **or**  $\frac{dy}{dx} = 2$  (o.e.) (c.a.o.) B1

Use of  $\text{grad}_{\text{normal}} \times \text{grad}_{\text{tangent}} = -1$  M1

Equation of normal:  $y - 1 = -\frac{1}{2}(x - 2)$   $\left[ \begin{array}{l} \text{f.t. candidate's value for } \frac{dy}{dx} \\ \frac{dy}{dx} \end{array} \right]$  A1

3. (a)  $8(2 \cos^2 \theta - 1) + 6 = \cos^2 \theta + \cos \theta$  (correct use of  $\cos 2\theta = 2 \cos^2 \theta - 1$ ) M1

An attempt to collect terms, form and solve quadratic equation

in  $\cos \theta$ , either by using the quadratic formula or by getting the

expression into the form  $(a \cos \theta + b)(c \cos \theta + d)$ ,

with  $a \times c =$  candidate's coefficient of  $\cos^2 \theta$  and  $b \times d =$  candidate's constant m1

$$15 \cos^2 \theta - \cos \theta - 2 = 0 \Rightarrow (5 \cos \theta - 2)(3 \cos \theta + 1) = 0$$

$$\Rightarrow \cos \theta = \frac{2}{5}, \cos \theta = -\frac{1}{3} \quad (\text{c.a.o.}) \quad \text{A1}$$

$$\theta = 66.42^\circ, 293.58^\circ \quad \text{B1}$$

$$\theta = 109.47^\circ, 250.53^\circ \quad \text{B1 B1}$$

Note: Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.

$\cos \theta = +, -, \text{f.t. for 3 marks, } \cos \theta = -, -, \text{f.t. for 2 marks}$

$\cos \theta = +, +, \text{f.t. for 1 mark}$

- (b)  $R = 4$  B1  
 Correctly expanding  $\cos(\theta + \alpha)$ , correctly comparing coefficients and using either  $4 \cos \alpha = \sqrt{15}$  or  $4 \sin \alpha = 1$  or  $\tan \alpha = \frac{1}{\sqrt{15}}$  to find  $\alpha$   $\sqrt{15}$   
 (f.t. candidate's value for  $R$ ) M1  
 $\alpha = 14.48^\circ$  (c.a.o.) A1  
 $\cos(\theta + 14.48^\circ) = \frac{3}{4} = 0.75$   
 (f.t. candidate's values for  $R, \alpha, 0^\circ < \alpha < 90^\circ$ ) B1  
 $\theta + 14.48^\circ = 41.41^\circ, 318.59^\circ$   
 (at least one value, f.t. candidate's values for  $R, \alpha, 0^\circ < \alpha < 90^\circ$ ) B1  
 $\theta = 26.93^\circ, 304.11^\circ$  (c.a.o.) B1

4.

$$\text{Volume} = \pi \int_{\pi/6}^{\pi/2} \sin^2 2x \, dx \quad \text{B1}$$

$$\sin^2 2x = \frac{(1 - \cos 4x)}{2} \quad \text{B1}$$

$$\int (a + b \cos 4x) \, dx = ax + \frac{1}{4} b \sin 4x, \quad a \neq 0, b \neq 0 \quad \text{B1}$$

Correct substitution of candidate's limits in candidate's integrated expression of form  $mx + n \sin 4x$   $m \neq 0, n \neq 0$  M1  
 Volume = 1.985 (c.a.o.) A1

**Note: Answer only with no working earns 0 marks**

5. (a) (i)  $(1 + 6x)^{1/3} = 1 + 2x - 4x^2$   $(1 + 2x)$  B1  
 $(-4x^2)$  B1  
 (ii)  $|x| < 1/6$  or  $-1/6 < x < 1/6$  B1
- (b)  $2 + 4x - 8x^2 = 2x^2 - 15x \Rightarrow 10x^2 - 19x - 2 = 0$  M1  
 (An attempt to set up and use a correct method to solve quadratic using candidate's expansion for  $(1 + 6x)^{1/3}$ )  
 $x = -0.1$  (f.t. only candidate's range for  $x$  in (a)) A1

6. (a) candidate's  $x$ -derivative =  $a$   
 candidate's  $y$ -derivative =  $-\frac{b}{t^2}$  (at least one term correct) B1
- $$\frac{dy}{dx} = \frac{\text{candidate's } y\text{-derivative}}{\text{candidate's } x\text{-derivative}} \quad \text{M1}$$
- $$\frac{dy}{dx} = -\frac{b}{at^2} \quad (\text{c.a.o.}) \quad \text{A1}$$
- Tangent at  $P$ :  $y - \frac{b}{p} = -\frac{b}{ap^2}(x - ap)$  (o.e.)
- (f.t. candidate's expression for  $\frac{dy}{dx}$ ) M1
- $$ap^2y - abp = -bx + abp$$
- $$ap^2y + bx - 2abp = 0. \quad (\text{convincing}) \quad \text{A1}$$
- (b)  $y = 0 \Rightarrow x = 2ap$  (o.e.) B1  
 $x = 0 \Rightarrow y = 2b/p$  (o.e.) B1  
 Area of triangle  $AOB = 2ab$  (c.a.o.) B1
- (c)  $p^2 - 2p + 2 = 0$  ( $abp^2 - 2abp + 2ab = 0$ ) B1  
 Attempting **either** to use the formula to solve the candidate's quadratic in  $p$  **or** to find the discriminant of the candidate's quadratic **or** to complete the square M1
- Either** discriminant  $< 0 \Rightarrow$  no real roots  $\Rightarrow$  no such  $P$  can exist **or**  $(p - 1)^2 + 1 = 0 \Rightarrow (p - 1)^2 = -1 \Rightarrow$  no such  $P$  can exist
- (c.a.o.) A1
7. (a)  $u = 3x - 1 \Rightarrow du = 3dx$  (o.e.) B1  
 $dv = \cos 2x \, dx \Rightarrow v = \frac{1}{2} \sin 2x$  (o.e.) B1
- $$\int (3x - 1) \cos 2x \, dx = (3x - 1) \times \frac{1}{2} \sin 2x - \int \frac{1}{2} \sin 2x \times 3 \, dx \quad \text{M1}$$
- $$\int (3x - 1) \cos 2x \, dx = \frac{1}{2} (3x - 1) \sin 2x + \frac{3}{4} \cos 2x + c \quad (\text{c.a.o.}) \quad \text{A1}$$

$$(b) \quad \int \frac{x}{(2x+1)^3} dx = \int \frac{f(u)}{u^3} \times k du \quad (f(u) = pu + q, p \neq 0, q \neq 0 \text{ and } k = \frac{1}{2} \text{ or } 2) \quad \text{M1}$$

$$\int \frac{x}{(2x+1)^3} dx = \int \frac{(u-1)}{2} \times \frac{1}{u^3} \times \frac{du}{2} \quad \text{A1}$$

$$\int (au^{-2} + bu^{-3}) du = \frac{au^{-1}}{-1} + \frac{bu^{-2}}{-2} \quad (a \neq 0, b \neq 0) \quad \text{B1}$$

**Either:** Correctly inserting limits of 1, 3 in candidate's  $cu^{-1} + du^{-2}$   
( $c \neq 0, d \neq 0$ )

**or:** Correctly inserting limits of 0, 1 in candidate's  
 $c(2x+1)^{-1} + d(2x+1)^{-2}$  ( $c \neq 0, d \neq 0$ ) m1

$$\int_0^1 \frac{x}{(2x+1)^3} dx = \frac{1}{18} \quad (= 0.055 \dots) \quad (\text{c.a.o.}) \quad \text{A1}$$

**Note:** Answer only with no working earns 0 marks

8. (a)  $\frac{dA}{dt} = k\sqrt{A}$  B1

(b)  $\int \frac{dA}{\sqrt{A}} = \int k dt$  M1

$$\frac{A^{1/2}}{1/2} = kt + c \quad \text{A1}$$

Substituting 64 for  $A$  and 3 for  $t$  and 196 for  $A$  and 5.5 for  $t$  in candidate's derived equation m1

$$16 = 3k + c, 28 = 5.5k + c \quad (\text{both equations}) \quad (\text{c.a.o.}) \quad \text{A1}$$

Attempting to solve candidate's derived simultaneous linear equations in  $k$  and  $c$  m1

$$A = (2.4t + 0.8)^2 \quad (\text{o.e.}) \quad (\text{c.a.o.}) \quad \text{A1}$$

9. (a)  $\mathbf{AB} = 8\mathbf{i} - 4\mathbf{j} + 12\mathbf{k}$  B1

(b)  $\mathbf{OC} = -\mathbf{i} + 3\mathbf{j} - 7\mathbf{k} + \frac{3}{4}(8\mathbf{i} - 4\mathbf{j} + 12\mathbf{k})$  (o.e.) M1  
 $\mathbf{OC} = 5\mathbf{i} + 2\mathbf{k}$  A1

(c) (i) Use of  $\mathbf{OA} + \mu(-4\mathbf{i} + \mathbf{j} + 3\mathbf{k})$  on r.h.s. M1  
 $\mathbf{r} = -\mathbf{i} + 3\mathbf{j} - 7\mathbf{k} + \mu(-4\mathbf{i} + \mathbf{j} + 3\mathbf{k})$  (all correct) A1

(ii)  $-1 + \lambda \times (-4) = 7$   
(an equation in  $\lambda$  from one set of coefficients) M1

$$\lambda = -2 \quad \text{A1}$$

$$1 + (-2) \times 1 = -1$$

$$11 + (-2) \times 3 = 5 \quad (\text{both verifications}) \quad \text{A1}$$

An attempt to evaluate  $\mathbf{AB} \cdot (-4\mathbf{i} + \mathbf{j} + 3\mathbf{k})$  M1

$$\mathbf{AB} \cdot (-4\mathbf{i} + \mathbf{j} + 3\mathbf{k}) = 0 \quad (\text{convincing}) \quad \text{A1}$$

$B$  lies on  $L$ ,  $AB$  is perpendicular to  $L$  and thus  $B$  is the foot of the perpendicular from  $A$  to  $L$  (c.a.o.) A1

**10.** Assume that there is a real value of  $x$  such that

$$(5x - 3)^2 + 1 < (3x - 1)^2.$$

$$25x^2 - 30x + 9 + 1 < 9x^2 - 6x + 1 \Rightarrow 16x^2 - 24x + 9 < 0 \quad \text{B1}$$

$$(4x - 3)^2 < 0 \quad \text{B1}$$

This contradicts the fact that  $x$  is real and thus  $(5x - 3)^2 + 1 \geq (3x - 1)^2$ . B1

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
<b>1</b>	$\begin{aligned} S_n &= \sum_{r=1}^n (2r-1)^2 = \sum_{r=1}^n 4r^2 - \sum_{r=1}^n 4r + \sum_{r=1}^n 1 \\ &= \frac{4n(n+1)(2n+1)}{6} - \frac{4n(n+1)}{2} + n \\ &= \frac{n}{6}(8n^2 + 12n + 4 - 12n - 12 + 6) \\ &= \frac{4n^3}{3} - \frac{n}{3} \quad \text{cao} \end{aligned}$	<b>M1A1</b>  <b>A1A1</b>   <b>A1</b>   <b>A1</b>	M1A0 for 2 correct terms  Award A1 for 2 correct  FT line above if at least 2 terms present  Penalise 1 mark if $n$ used as dummy variable in summations
<b>2(a)</b>	<p>EITHER     <math>\frac{1}{w} = \frac{1}{1-i} + \frac{1}{1+2i}</math></p> $= \frac{1+2i+1-i}{(1-i)(1+2i)}$ $= \frac{2+i}{3+i}$ $w = \frac{3+i}{2+i} \times \frac{2-i}{2-i}$ $= \frac{7-i}{5}$ <p>OR            <math>\frac{1}{1-i} = \frac{1+i}{1-i^2} = \frac{1+i}{2}</math></p> $\frac{1}{1+2i} = \frac{1-2i}{1-4i^2} = \frac{1-2i}{5}$ $\frac{1}{w} = \frac{5+5i+2-4i}{10} = \frac{7+i}{10}$ $w = \frac{10}{7+i} \times \frac{7-i}{7-i}$ $= \frac{7-i}{5}$	  <b>M1A1</b>  <b>A1</b>   <b>M1</b>  <b>A1A1</b>   <b>M1A1</b>  <b>A1</b>  <b>A1</b>   <b>M1</b>  <b>A1</b>	        1 each for num and denom          1 each for num and denom
<b>(b)</b>	Mod( $w$ ) = $\frac{\sqrt{50}}{5}$ ( $\sqrt{2}$ ) Arg( $w$ ) = $-0.142$ ( $-8.13^\circ$ )	       <b>B1</b>  <b>B1</b>	       FT on their $w$ Accept $351.9^\circ$ or $6.14$ Do not FT arg if in 1 <sup>st</sup> quadrant

<p><b>3(a)</b></p>	$\alpha + \beta + \gamma = 2, \beta\gamma + \gamma\alpha + \alpha\beta = 2, \alpha\beta\gamma = -1$ $\frac{\beta\gamma}{\alpha} + \frac{\gamma\alpha}{\beta} + \frac{\alpha\beta}{\gamma} = \frac{\beta^2\gamma^2 + \gamma^2\alpha^2 + \alpha^2\beta^2}{\alpha\beta\gamma}$ $= \frac{(\beta\gamma + \gamma\alpha + \alpha\beta)^2 - 2\alpha\beta\gamma(\alpha + \beta + \gamma)}{\alpha\beta\gamma}$ $= \frac{(2)^2 - 2 \times (-1) \times 2}{-1} = -8$ <p><b>(b)</b> Consider</p> $\frac{\gamma\alpha}{\beta} \times \frac{\alpha\beta}{\gamma} + \frac{\alpha\beta}{\gamma} \times \frac{\beta\gamma}{\alpha} + \frac{\beta\gamma}{\alpha} \times \frac{\gamma\alpha}{\beta}$ $= \alpha^2 + \beta^2 + \gamma^2$ $= (\alpha + \beta + \gamma)^2 - 2(\beta\gamma + \gamma\alpha + \alpha\beta)$ $= 4 - 2 \times 2 = 0$ <p>Consider</p> $\frac{\beta\gamma}{\alpha} \times \frac{\gamma\alpha}{\beta} \times \frac{\alpha\beta}{\gamma} = \alpha\beta\gamma = -1$ <p>The required equation is</p> $x^3 + 8x^2 + 1 = 0$	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1A1</b></p> <p><b>B1</b></p>	<p>Convincing</p> <p>FT their coefficients</p>
--------------------	---	--	--




<p><b>4(a)</b></p>	<p>Rotation matrix = <math>\begin{bmatrix} 0 &amp; -1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix}</math></p> <p>Translation matrix = <math>\begin{bmatrix} 1 &amp; 0 &amp; 2 \\ 0 &amp; 1 &amp; 1 \\ 0 &amp; 0 &amp; 1 \end{bmatrix}</math></p> <p>Ref matrix in <math>y = x = \begin{bmatrix} 0 &amp; 1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix}</math></p> <p><math>T = \begin{bmatrix} 0 &amp; 1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \begin{bmatrix} 1 &amp; 0 &amp; 2 \\ 0 &amp; 1 &amp; 1 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \begin{bmatrix} 0 &amp; -1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} =</math></p> <p><math>\begin{bmatrix} 0 &amp; 1 &amp; 1 \\ 1 &amp; 0 &amp; 2 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \begin{bmatrix} 0 &amp; -1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \text{ or } \begin{bmatrix} 0 &amp; 1 &amp; 0 \\ 1 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \begin{bmatrix} 0 &amp; -1 &amp; 2 \\ 1 &amp; 0 &amp; 1 \\ 0 &amp; 0 &amp; 1 \end{bmatrix}</math></p> <p><math>= \begin{bmatrix} 1 &amp; 0 &amp; 1 \\ 0 &amp; -1 &amp; 2 \\ 0 &amp; 0 &amp; 1 \end{bmatrix}</math></p> <p><b>(b)</b> Fixed points satisfy</p> <p><math>\begin{bmatrix} 1 &amp; 0 &amp; 1 \\ 0 &amp; -1 &amp; 2 \\ 0 &amp; 0 &amp; 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}</math></p> <p><math>x = x + 1, (y = -y + 2)</math></p> <p>These equations are not consistent so there are no fixed points.</p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>Accept equivalent reason</p>
<p><b>5</b></p>	<p>Putting <math>n = 1</math>, the formula gives 6 which is divisible by 6 so the result is true for <math>n = 1</math></p> <p>Assume formula is true for <math>n = k</math>, ie</p> <p><math>7^k - 1</math> is divisible by 6 or <math>7^k = 6N + 1</math></p> <p>Consider, for <math>n = k + 1</math>,</p> <p><math>7^{k+1} - 1 = 7 \cdot 7^k - 1</math></p> <p><math>= 7(6N + 1) - 1</math></p> <p><math>= 42N + 6</math></p> <p>This is divisible by 6 therefore true for <math>n = k \Rightarrow</math> true for <math>n = k + 1</math> and since true for <math>n = 1</math>, the result is proved by induction.</p>	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	

<b>6(a)(i)</b>	$\text{Det}(\mathbf{A}) = 7 - 4\lambda + \lambda(5\lambda - 14) + 3(8 - 5)$ $= 5\lambda^2 - 18\lambda + 16$	<b>M1</b>	
<b>(ii)</b>	Putting $\lambda = 2$ , $\det = 20 - 36 + 16 = 0$ So $\mathbf{A}$ is singular. Putting $\det(\mathbf{A}) = 0$ , product of roots is $16/5$ So the other root is $8/5$	<b>A1</b> <b>B1</b>	
<b>(b)(i)</b>	$\begin{aligned} x + 2y + 3z &= 2 \\ 2x + y + 2z &= 1 \\ 5x + 4y + 7z &= 4 \end{aligned}$ <p>Attempting to use row operations,</p> $\begin{aligned} x + 2y + 3z &= 2 \\ 3y + 4z &= 3 \\ 6y + 8z &= 6 \end{aligned}$ <p>Since the 3<sup>rd</sup> equation is twice the 2<sup>nd</sup> equation, it follows that the equations are consistent.</p>	<b>M1</b> <b>A1</b> <b>A1</b>	
<b>(ii)</b>	Let $z = \alpha$ $y = 1 - \frac{4}{3}\alpha$ $x = -\frac{1}{3}\alpha$ (or equivalent)	<b>M1</b> <b>A1</b> <b>A1</b>	Or because the next step gives a row of zeroes
<b>(c)(i)</b>	$\mathbf{A} = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 1 & 1 \\ 5 & 4 & 7 \end{bmatrix}$ $\text{Cofactor matrix} = \begin{bmatrix} 3 & -9 & 3 \\ 5 & -8 & 1 \\ -2 & 5 & -1 \end{bmatrix} \text{ si}$ $\text{Adjugate matrix} = \begin{bmatrix} 3 & 5 & -2 \\ -9 & -8 & 5 \\ 3 & 1 & -1 \end{bmatrix}$	<b>M1A1</b>	Award M1 if at least 5 correct elements
<b>(ii)</b>	Determinant = 3 $\text{Inverse matrix} = \frac{1}{3} \begin{bmatrix} 3 & 5 & -2 \\ -9 & -8 & 5 \\ 3 & 1 & -1 \end{bmatrix}$	<b>A1</b> <b>B1</b> <b>B1</b>	No FT from incorrect cofactor matrix  FT from incorrect adjugate
<b>(iii)</b>	$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 3 & 5 & -2 \\ -9 & -8 & 5 \\ 3 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}$	<b>M1</b> <b>A1</b>	FT from inverse matrix

	$= \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$		
7	<p>Taking logs,</p> $\ln f(x) = \ln \sqrt{1 + \sin x} - \ln(1 + \tan x)^2$ $= \frac{1}{2} \ln(1 + \sin x) - 2 \ln(1 + \tan x)$ <p>Differentiating,</p> $\frac{f'(x)}{f(x)} = \frac{\cos x}{2(1 + \sin x)} - \frac{2 \sec^2 x}{(1 + \tan x)}$ <p>Putting <math>x = \pi/4</math>,</p> $f'(\pi/4) = -0.586 \text{ cao}$	<p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>B3</b></p> <p><b>M1</b></p> <p><b>A2</b></p>	<p>B1 for each correct term</p>
8(a)	$u + iv = (x + iy)^2$ $= x^2 - y^2 + 2ixy$ <p>Equating real and imaginary parts,</p> $u = x^2 - y^2$ $v = 2xy$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	
(b)	<p>Substituting for y,</p> $u = x^2 - (2x^2 + 1) = -1 - x^2$ $v^2 = 4x^2(2x^2 + 1)$ <p>Eliminating x,</p> $x^2 = -(u + 1)$ <p>So that</p> $v^2 = 4(u + 1)(2u + 1) \text{ cao}$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>FT their expressions from (a)</p>


## FP2

Ques	Solution	Mark	Notes
1	$u = x^2 \Rightarrow du = 2x dx,$ $[1, 2] \rightarrow [1, 4]$ $I = \frac{1}{2} \int_1^4 \frac{du}{\sqrt{25 - u^2}}$ $= \frac{1}{2} \left[ \sin^{-1} \left( \frac{u}{5} \right) \right]_1^4$ $= 0.363 \text{ cao}$	<b>B1</b> <b>B1</b>  <b>M1</b>   <b>A1</b>  <b>A1</b>	
2(a)	Substituting $t = \tan(\theta/2)$ $\frac{2t}{1+t^2} + \frac{3(1-t^2)}{1+t^2} = 2$ $2t + 3 - 3t^2 = 2 + 2t^2$ $5t^2 - 2t - 1 = 0$ $t = \frac{2 \pm \sqrt{24}}{10} = 0.68989\dots, -0.28989\dots$	<b>M1A1</b>   <b>A1</b>  <b>M1A1</b>	Convincing.  FT their roots from (a)
(b)	$t = 0.68989\dots$ giving $\theta/2 = 0.6039\dots$ The general solution is $\theta = 1.21 + 2n\pi$ $t = -0.28989\dots$ giving $\theta/2 = -0.2821\dots$ The general solution is $\theta = -0.564 + 2n\pi$	<b>B1</b> <b>B1</b> <b>B1</b> <b>B1</b>	Accept 2.859... Accept $5.72 + 2n\pi$

<p><b>3(a)</b></p>	$-1 = \cos \pi + i \sin \pi$ $\sqrt[4]{-1} = \cos \pi/4 + i \sin \pi/4 = \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}}$ $\text{Root2} = \cos 3\pi/4 + i \sin 3\pi/4 = -\frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}}$ $\text{Root3} = \cos 5\pi/4 + i \sin 5\pi/4 = -\frac{1}{\sqrt{2}} - i \frac{1}{\sqrt{2}}$ $\text{Root4} = \cos 7\pi/4 + i \sin 7\pi/4 = \frac{1}{\sqrt{2}} - i \frac{1}{\sqrt{2}}$	<p><b>B1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>Special case : Award 2/6 if they misread <math>-1</math> as <math>1</math>.</p>
<p><b>(b)(i)</b></p>		<p><b>B1</b></p>	<p>FT their roots if possible</p>
<p><b>(ii)</b></p>	<p>Length of side = <math>\frac{2}{\sqrt{2}}</math></p> <p>Area of square = 2</p>	<p><b>B1</b></p> <p><b>B1</b></p>	

<p><b>4(a)</b></p>	$f'(x) = \frac{2(x-1) - (2x+3)}{(x-1)^2}$ $= -\frac{5}{(x-1)^2}$ <p>This is negative for all <math>x &gt; 1</math> therefore <math>f</math> is strictly decreasing.</p>	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	
<p><b>(b)(i)</b></p> <p><b>(ii)</b></p>	<p><math>f(4) = 11/3, f(5) = 13/4</math>  <math>f(S) = [13/4, 11/3]</math></p> <p>EITHER</p> $y = \frac{2x+3}{x-1} \Rightarrow x = \frac{y+3}{y-2}$ <p><math>f^{-1}(4) = 7/2, f^{-1}(5) = 8/3</math>  <math>f^{-1}(S) = [8/3, 7/2]</math></p> <p>OR</p> $\frac{2x+3}{x-1} = 4 \rightarrow x = \frac{7}{2}$ $\frac{2x+3}{x-1} = 5 \rightarrow x = \frac{8}{3}$ <p><math>f^{-1}(S) = [8/3, 7/2]</math></p>	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>A0 if wrong way around but penalise only once.</p> <p>A0 if wrong way around.</p> <p>M1A1 for the first and then A1 for the second.</p> <p>A0 if wrong way around.</p>
<p><b>5(a)(i)</b></p> <p><b>(ii)</b></p> <p><b>(iii)</b></p> <p><b>(iv)</b></p> <p><b>(b)(i)</b></p> <p><b>(ii)</b></p>	<p>Completing the square,  <math>(x-2)^2 + 2(y+1)^2 = 4</math>  The centre is therefore <math>(2, -1)</math>  In standard form, the equation is  <math>\frac{(x-2)^2}{4} + \frac{(y+1)^2}{2} = 1</math> so <math>a = 2, b = \sqrt{2}</math> si  <math display="block">e = \sqrt{\frac{4-2}{4}} = \frac{1}{\sqrt{2}}</math>  The foci are <math>(2 + \sqrt{2}, -1)</math> and <math>(2 - \sqrt{2}, -1)</math></p> <p>The equations of the directrices are <math>x = 2 \pm 2\sqrt{2}</math></p> <p>EITHER</p> <p>Putting <math>x = 0, (y+1)^2 = 0</math>  This has a repeated root, hence <math>x = 0</math> is a tangent  OR  Semi-major axis = 2 = <math>x</math>-coordinate of centre  This equality shows that <math>x = 0</math> is a tangent  Substituting <math>y = mx</math>,  <math>x^2(1 + 2m^2) - x(4 - 4m) + 2 = 0</math>  Use of the condition for tangency, ie '<math>b^2 = 4ac</math>'  <math>16(1 - m)^2 = 8(1 + 2m^2)</math>  <math>2 - 4m + 2m^2 = 1 + 2m^2 \Rightarrow m = \frac{1}{4}</math></p>	<p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>M1A1</b></p> <p><b>B1B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>FT their equation in (ii), (iii) and (iv)</p>

<p><b>6(a)</b></p> <p><b>(b)</b></p>	<p>Let</p> $\frac{4x^2 - 2x + 9}{x(x^2 + 3)} \equiv \frac{A}{x} + \frac{Bx + C}{x^2 + 3}$ $= \frac{A(x^2 + 3) + x(Bx + C)}{x(x^2 + 3)} \quad (\text{oe})$ <p><math>x = 0</math> gives <math>A = 3</math></p> <p>Coeff of <math>x^2</math> gives <math>A + B = 4</math>, <math>B = 1</math></p> <p>Coeff of <math>x</math> gives <math>C = -2</math></p> $\int_1^3 \frac{4x^2 - 2x + 9}{x(x^2 + 3)} dx = \int_1^3 \left( \frac{3}{x} + \frac{x}{x^2 + 3} - \frac{2}{x^2 + 3} \right) dx$ $= \left[ 3 \ln x + \frac{1}{2} \ln(x^2 + 3) - \frac{2}{\sqrt{3}} \tan^{-1} \left( \frac{x}{\sqrt{3}} \right) \right]_1^3$ $= 3 \ln 3 + \frac{1}{2} \ln 12 - \frac{2}{\sqrt{3}} \tan^{-1} \left( \frac{3}{\sqrt{3}} \right)$ $- 3 \ln 1 - \frac{1}{2} \ln 4 + \frac{2}{\sqrt{3}} \tan^{-1} \left( \frac{1}{\sqrt{3}} \right)$ $= 3.24 \quad \text{cao}$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>B3</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>B1 each term</p>
--------------------------------------	---	---	---------------------

<p><b>7(a)</b></p>	<p>Consider</p> $f(-x) = \frac{(2(-x)^2 + 1)^2}{(-x)^3} = -f(x)$ <p>Therefore <math>f</math> is odd</p>	<p><b>M1A1</b></p> <p><b>A1</b></p>	
<p><b>(b)</b></p>	<p>EITHER</p> <p>Differentiating,</p> $f'(x) = \frac{2(2x^2 + 1).4x.x^3 - 3x^2(2x^2 + 1)^2}{x^6}$ <p>At a stationary point, putting <math>f'(x) = 0</math>,</p> $8x^2 = 3(2x^2 + 1)$ $x = \pm\sqrt{\frac{3}{2}}$ <p>OR</p> <p>Consider <math>f(x) = 4x + \frac{4}{x} + \frac{1}{x^3}</math></p> $f'(x) = 4 - \frac{4}{x^2} - \frac{3}{x^4}$ <p>At a stationary point, putting <math>f'(x) = 0</math>,</p> $4x^4 - 4x^2 - 3 = 0$ $x = \pm\sqrt{\frac{3}{2}}$	<p><b>M1A1</b></p> <p><b>m1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>m1</b></p> <p><b>A1</b></p>	<p>Condone the cancellation of <math>x^2(2x^2 + 1)</math></p>
<p><b>(c)</b></p>	<p>The asymptotes are</p> $x = 0$ $y = 4x$	<p><b>B1</b></p> <p><b>B1</b></p>	
<p><b>(d)</b></p>		<p><b>G1</b></p> <p><b>G1</b></p>	



8	<p>EITHER</p> <p>Consider</p> $\cos 5\theta + i\sin 5\theta = (\cos \theta + i\sin \theta)^5$ <p>Expanding and taking real parts,</p> $\cos 5\theta = \cos^5 \theta + 10\cos^3 \theta (i\sin \theta)^2 + 5\cos \theta (i\sin \theta)^4$ $= \cos^5 \theta - 10\cos^3 \theta (1 - \cos^2 \theta) + 5\cos \theta (1 - \cos^2 \theta)^2$ $= \cos^5 \theta - 10\cos^3 \theta + 10\cos^5 \theta + 5\cos \theta - 10\cos^3 \theta + 5\cos^5 \theta$ $= 16\cos^5 \theta - 20\cos^3 \theta + 5\cos \theta$ <p>OR</p> <p>Let <math>z = \cos \theta + i\sin \theta</math></p> <p>So that <math>z + \frac{1}{z} = 2\cos \theta</math> and <math>z^n + \frac{1}{z^n} = 2\cos n\theta</math></p> <p>Consider</p> $\left(z + \frac{1}{z}\right)^5 = z^5 + 5z^3 + 10z + \frac{10}{z} + \frac{5}{z^3} + \frac{1}{z^5}$ $32\cos^5 \theta = 2\cos 5\theta + 10\cos 3\theta + 20\cos \theta$ $\cos 5\theta = 16\cos^5 \theta - 5\cos 3\theta - 10\cos \theta$ $= 16\cos^5 \theta - 5(4\cos^3 \theta - 3\cos \theta) - 10\cos \theta$ $= 16\cos^5 \theta - 20\cos^3 \theta + 5\cos \theta$	<p><b>M1</b></p> <p><b>m1A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	
---	---	--	--

**FP3**

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
<b>1</b>	Using $\cosh 2x = 2\cosh^2 x - 1$ , the eqn becomes $2\cosh^2 x - 7\cosh x + 6 = 0$ Solving the quadratic equation, $\cosh x = 2, 1.5$ The positive roots are therefore $x = \cosh^{-1} 2 = 1.32$ and $x = \cosh^{-1}(1.5) = 0.96$	M1 A1 M1 A1  A1 A1	FT their roots
<b>2(a)(i)</b>	The Newton-Raphson iteration is $x_{n+1} = x_n - \frac{(x_n^3 - a)}{3x_n^2}$ $= \frac{2x_n^3 + a}{3x_n^2}$	M1  A1	Convincing
<b>(ii)</b>	$x_0 = 2$ $x_1 = 2.166666667$ $x_2 = 2.154503616$ $x_3 = 2.154434692$ $x_4 = 2.15443469$ $\sqrt[3]{10} = 2.1544$ correct to 4 decimal places.	M1A1     A1	
<b>(b)</b>	Consider $\frac{d}{dx} \left( \frac{a}{x^2} \right) = -\frac{2a}{x^3}$ $= -2 \text{ when } x = \sqrt[3]{a}$ The sequence diverges because this exceeds 1 in modulus.	M1A1  A1  A1	M0 if $a = 10$
<b>3(a)</b>	$f'(x) = \frac{2e^x}{2e^x - 1}$	B1	
	$f''(x) = \frac{2e^x(2e^x - 1) - 2e^x \cdot 2e^x}{(2e^x - 1)^2}$	M1	
	$= \frac{-2e^x}{(2e^x - 1)^2}$	A1	convincing
<b>(b)</b>	$f'''(x) = \frac{-2e^x(2e^x - 1)^2 + 2e^x \cdot 2e^x \cdot 2(2e^x - 1)}{(2e^x - 1)^4}$	M1A1	
	$f(0) = 0, f'(0) = 2, f''(0) = -2, f'''(0) = 6$ The Maclaurin series is $2x - x^2 + x^3 + \dots$	B2  M1A1	Award B1 for 2 correct values  FT on their values of $f^{(n)}(0)$

4	<p>Completing the square,</p> $3 + 2x - x^2 = 4 - (x-1)^2$ <p>so <math>I = \int_1^2 \sqrt{4 - (x-1)^2} dx</math></p> <p>Put <math>x-1 = 2\sin\theta</math>  <math>dx = 2\cos\theta d\theta, [1,2] \rightarrow [0,\pi/6]</math></p> $I = \int_0^{\pi/6} \sqrt{4 - 4\sin^2\theta} \cdot 2\cos\theta d\theta$ $= 4 \int_0^{\pi/6} \cos^2\theta d\theta$ $= 2 \int_0^{\pi/6} (1 + \cos 2\theta) d\theta$ $= 2 \left[ \theta + \frac{\sin 2\theta}{2} \right]_0^{\pi/6}$ $= 1.91$	<p>M1A1</p> <p>M1 A1A1</p> <p>m1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>Allow <math>x-1 = 2\cos\theta</math></p>
<p>5(a)</p> <p>(b)</p>	$I_n = \left[ x^n \cosh x \right]_0^1 - n \int_0^1 x^{n-1} \cosh x dx$ $= \cosh 1 - n \int_0^1 x^{n-1} \cosh x dx$ $= \cosh 1 - \left[ nx^{n-1} \sinh x \right]_0^1 + n(n-1)I_{n-2}$ $= \cosh 1 - n \sinh 1 + n(n-1)I_{n-2}$ $I_0 = \int_0^1 \sinh x dx = \left[ \cosh x \right]_0^1 = \cosh 1 - 1$ $I_4 = \cosh 1 - 4 \sinh 1 + 12I_2$ $= \cosh 1 - 4 \sinh 1 + 12(\cosh 1 - 2 \sinh 1 + 2I_0)$ $= 13 \cosh 1 - 28 \sinh 1 + 24(\cosh 1 - 1)$ $= 37 \cosh 1 - 28 \sinh 1 - 24 \quad \text{cao}$	<p>M1A1</p> <p>A1</p> <p>M1A1</p> <p>M1A1</p> <p>M1 A1</p> <p>A1</p>	<p>M1A1 for evaluating <math>I_0</math> at any stage</p> <p>FT their <math>I_0</math> if substituted here</p>

<p><b>6(a)</b></p>	<p>Consider</p> $x = r \cos \theta$ $= \sin^2 \theta \cos \theta$ $\frac{dx}{d\theta} = 2 \sin \theta \cos^2 \theta - \sin^3 \theta$ <p>The tangent is perpendicular to the initial line where <math>\frac{dx}{d\theta} = 2 \sin \theta \cos^2 \theta - \sin^3 \theta = 0</math></p> $\tan^2 \theta = 2$ $\theta = \tan^{-1} \sqrt{2} = 0.955$ $r = 0.667$	<p>M1 A1 M1A1</p>	<p>Do not penalise the removal of the factor <math>\sin \theta</math></p>
<p><b>(b)</b></p>	<p>Area = <math>\frac{1}{2} \int r^2 d\theta</math></p> $= \frac{1}{2} \int_0^{\pi/2} (1 - \sin \theta)^2 d\theta$ $= \frac{1}{2} \int_0^{\pi/2} (1 - 2 \sin \theta + \sin^2 \theta) d\theta$ $= \frac{1}{4} \int_0^{\pi/2} (3 - 4 \sin \theta - \cos 2\theta) d\theta$ $= \frac{1}{4} \left[ 3\theta + 4 \cos \theta - \frac{1}{2} \sin 2\theta \right]_0^{\pi/2}$ $= \frac{3\pi - 8}{8} \quad (0.178) \quad \text{cao}$	<p>M1 A1 A1 A1 A1 A1</p>	

7(a)(i)	$D(\operatorname{cosech} x) = D\left(\frac{1}{\sinh x}\right)$ $= \frac{-1}{\sinh^2 x} \times \cosh x$ $= -\operatorname{cosech} x \coth x$ $D(\coth x) = D\left(\frac{\cosh x}{\sinh x}\right)$ $= \frac{\sinh^2 x - \cosh^2 x}{\sinh^2 x}$ $= -\operatorname{cosech}^2 x$	M1 A1	convincing
(ii)	$D \ln(\operatorname{cosech} x + \coth x)$ $= \frac{-(\operatorname{cosech} x \coth x + \operatorname{cosech}^2 x)}{(\operatorname{cosech} x + \coth x)}$ $= -\operatorname{cosech} x$	M1 A1	
(b)(i)	$L = \int \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ $= \int_1^e \sqrt{1 + \left(\frac{1}{x}\right)^2} dx$ $= \int_1^e \frac{\sqrt{1+x^2}}{x} dx$	M1 A1	
(ii)	<p>Putting <math>x = \sinh u</math>,  <math>dx = \cosh u du</math>, <math>[1, e] \rightarrow [\sinh^{-1} 1, \sinh^{-1} e]</math> (<math>[\alpha, \beta]</math>)</p> $\text{Arc length} = \int_{\alpha}^{\beta} \frac{\sqrt{1 + \sinh^2 u}}{\sinh u} \cdot \cosh u du$ $= \int_{\alpha}^{\beta} \frac{\cosh^2 u}{\sinh u} du$ $= \int_{\alpha}^{\beta} \frac{1 + \sinh^2 u}{\sinh u} du$ $= \int_{\alpha}^{\beta} (\operatorname{cosech} u + \sinh u) du$	B1B1 M1 A1 A1	
(iii)	$= [-\ln(\operatorname{cosech} u + \coth u) + \cosh u]_{\alpha}^{\beta}$ $= 2.00$	M1A1 A2	



WJEC  
245 Western Avenue  
Cardiff CF5 2YX  
Tel No 029 2026 5000  
Fax 029 2057 5994  
E-mail: [exams@wjec.co.uk](mailto:exams@wjec.co.uk)  
website: [www.wjec.co.uk](http://www.wjec.co.uk)



# **GCE MARKING SCHEME**

## **MATHEMATICS - M1-M3 & S1-S3 AS/Advanced**

**SUMMER 2013**

## INTRODUCTION

The marking schemes which follow were those used by WJEC for the Summer 2013 examination in GCE MATHEMATICS. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

<b>Paper</b>	<b>Page</b>
M1	1
M2	9
M3	17
S1	23
S2	26
S3	29




# M1

Q

Solution

Mark Notes



1(a)

B1 (0, 20) to (8, 20)  
Or (18, 6) to (40, 6)  
B1 (8, 20) to (18, 6)  
B1 completely correct with all units and labels.

1(b) Deceleration = gradient of graph

$$D = \frac{20-6}{18-8}$$

$$D = \underline{1.4 \text{ ms}^{-2}}$$

M1 any correct method

A1 ft graph +/-

A1 cao

OR

Use of  $v = u + at$ ,  $v=6$ ,  $u=20$ ,  $t=10$

$$6 = 20 + 10a$$

$$a = -1.4 \text{ ms}^{-2}$$

Magnitude of acceleration =  $1.4 \text{ ms}^{-2}$

M1

A1 allow  $-a$


A1 cao

1(c) Distance AB = Area under graph  
 $= (8 \times 20) + 0.5(20 + 6) \times 10 + (22 \times 6)$   
 $= 160 + 130 + 132$   
 $= \underline{422 \text{ m}}$

M1 used. Oe

B1 any correct area, ft graph

A1 cao



2(a)

N2L applied to lift and person


$$(M + 64)g - 7500 = (M+64) \times 0.425$$

$$M = \underline{736}$$

M1 dim correct equation,  
forces opposing

A1 correct equation

A1



2(b)

N2L applied to person

$$64g - R = 64a$$

$$R = 64 \times 9.8 - 64 \times 0.425$$

$$R = \underline{600 \text{ N}}$$


M1 64g and R opposing  
Dim correct equation

A1 correct equation

A1

Q	Solution	Mark	Notes
3(a)	$v^2 = u^2 + 2as$ , $v=0$ , $a=(\pm)9.8$ , $s=18.225$ $0 = u^2 - 2 \times 9.8 \times 18.225$ $u = \underline{18.9}$	M1 A1 A1	oe used  convincing
3(b)	Use of $s = ut + 0.5at^2$ , $s=(\pm)2.8$ , $a=(\pm)9.8$ , $u=18.9$ $-2.8 = 18.9t + 0.5 \times (-9.8)t^2$ $4.9t^2 - 18.9t - 2.8 = 0$ $7t^2 - 27t - 4 = 0$ $(7t + 1)(t - 4) = 0$  $t = \underline{4s}$	M1 A1    m1 A1	oe     correct method for solving quad equ seen cao


Q	Solution	Mark	Notes
---	----------	------	-------



4

5  
~

4(a)	N2L applied to B $5g - T = 5a$	M1	dim correct equation 5g and T opposing.
	$T = 5 \times 9.8 - 5 \times 1.61$	A1	
	$T = \underline{40.95 \text{ N}}$	A1	cao
	$R = 9g = (88.2 \text{ N})$	B1	si
	$F = 9\mu g = (88.2\mu)$	B1	si
	N2L applied to A	M1	dim correct equation T and F opposing
	$T - F = 9a$	A1	
	$T - 88.2\mu = 9 \times 1.61$		
	$\mu = \underline{0.3}$	A1	cao
4(b)	limiting friction $= 9\mu g = 9 \times 0.6g = 5.4g$	B1	
	Limiting friction $> 5g$		
	Particle will remain at rest	R1	oe
	$T = 5g = \underline{49 \text{ N}}$	B1	



5

5(a)(i) Resolve vertically

$$R + 84 = 12g$$

$$R = \underline{33.6}$$

M1 all forces, no extras

A1

A1 cao

5(a)(ii) Moments about C

$$12g \times 0.2 = 84(x - 0.8)$$

$$84x = 12g \times 0.2 + 84 \times 0.8$$

$$x = \underline{1.08}$$

M1 equation, no extra force  
oe

B1 any correct moment

A1 correct equation

A1 cao

5(b) When about to tilt about C,  $R_D = 0$ 

Moments about C


$$Mg \times 0.8 = 12g \times 0.2$$

$$M = \underline{3}$$

M1 si

m1 equation, no extra force

A1

Q	Solution	Mark	Notes
<hr/>			
			
6.			
6(a)	<p>Conservation of momentum</p> $2u + 5 \times 0 = 2 \times (-2) + 5 \times 3$ $u = \underline{5.5}$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>equation required, only 1 sign error.</p> <p>correct equation</p>
6(b)	<p>Restitution</p> $3 - (-2) = -e(0 - 5.5)$ $e = \frac{10}{11} = 0.909$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>only 1 sign error</p> <p>ft <math>u</math></p> <p>cao</p>
6(c)	<p>Impulse = change of momentum</p> $I = 5(3 - 0)$ $I = \underline{15 \text{ (Ns)}}$	<p>M1</p> <p>A1</p>	<p>for <math>P</math> or <math>Q</math></p> <p>+ required</p>
6(d)	$v' = ev$ $v' = 0.25 \times 3$ $v' = \underline{0.75 \text{ ms}^{-1}}$	<p>M1</p> <p>A1</p>	<p>used</p> <p>+ required</p>

Q	Solution	Mark	Notes
7.(a)	Resolve $X = 85 - 40 + 75 \cos \alpha$ $X = 85 - 40 + 75 \times 0.8$ $X = 105$	M1 B1 A1	attempted any correct resolution all correct accept $\cos 36.9$
	Resolve $Y = 60 - 75 \sin \alpha$ $Y = 60 - 75 \times 0.6$ $Y = 15$	M1 A1	attempted all correct, accept $\sin 36.9$
	$R = \sqrt{105^2 + 15^2}$ $R = 75\sqrt{2} = \underline{106.066 \text{ N}}$	M1 A1	 cao
	$\theta = \tan^{-1}\left(\frac{15}{105}\right)$ $\theta = \underline{8.13^\circ}$	M1 A1	allow reciprocal cao
7(b)	N2L applied to particle $75\sqrt{2} = 5a$ $a = 15\sqrt{2} = \underline{21.21 \text{ ms}^{-2}}$	M1 A1	dim correct equation ft $R$ if first 2 M's gained.

Q	Solution			Mark	Notes
8.	Area	from $AD$	from $AB$		
	$APCD$ 48	3	4	B1	
	$PBC$ 24	8	$8/3$	B1	
	Circle $4\pi$	3	3	B1	
	Lamina $(72-4\pi)$	$x$	$y$	B1	areas
8(a)	Moments about $AD$			M1	equation
	$48 \times 3 + 24 \times 8 = 4\pi \times 3 + (72 - 4\pi)x$			A1	ft table
	$x = \underline{5.02 \text{ cm}}$			A1	cao
	Moments about $AB$			M1	equation
	$48 \times 4 + 24 \times 8/3 = 4\pi \times 3 + (72 - 4\pi)y$			A1	ft table
	$y = \underline{3.67 \text{ cm}}$			A1	cao
8(b)	$AQ = \underline{3.67 \text{ cm}}$			B1	ft y



## M2

Q	Solution	Mark	Notes
1(a)	$\begin{aligned}\text{Loss in KE} &= 0.5mv^2 \\ &= 0.5 \times 8 \times 7^2 \\ &= \underline{196\text{J}}\end{aligned}$	M1 A1	Corr use of KE formula
1(b)	<p>Work energy principle</p> $196 = F \times 15$ $F = \mu R$ $= 8g\mu = (78.4\mu)$ <p>Therefore <math>196 = 78.4\mu \times 15</math></p> $\mu = \frac{1}{6}$	M1 A1  B1	correct use ft loss in KE
	<p>OR</p> <p>Use of <math>v^2 = u^2 + 2as</math></p> $0 = 7^2 + 2a \times 15$ $a = -1.633$	(M1)	
	<p>Use <math>F = ma</math></p> $-F = 8 \times -1.633$ $F = 8\mu g$ $\mu = \frac{13.067}{8g} = \frac{1}{6}$	(M1) (B1) (A1)p	

Q	Solution	Mark	Notes
2(a)	$\mathbf{r} = \int \mathbf{v} dt$ $\mathbf{r} = \int (13t - 3)\mathbf{i} + (2 + 3t^2)\mathbf{j} dt$ $\mathbf{r} = \left(\frac{13}{2}t^2 - 3t\right)\mathbf{i} + (2t + t^3)\mathbf{j} + (\underline{c})$	M1  A1 A1	use of integration  one for each coefficient
	When $t = 0$ , $\mathbf{c} = 2\mathbf{i} + 7\mathbf{j}$ $\mathbf{r} = (6.5t^2 - 3t + 2)\mathbf{i} + (2t + t^3 + 7)\mathbf{j}$	m1 A1	use of initial conditions ft $\mathbf{r}$
2(b)	$\mathbf{a} = \frac{d\mathbf{v}}{dt}$ $= 13\mathbf{i} + 6t\mathbf{j}$	 A1	 M1 use of differentiation
2(c)	We require $\mathbf{v} \cdot (\mathbf{i} - 2\mathbf{j}) = 0$ $\mathbf{v} \cdot (\mathbf{i} - 2\mathbf{j}) = (13t - 3) - 2(2 + 3t^2)$ $= -6t^2 + 13t - 7$ $6t^2 - 13t + 7 = 0$ $(6t - 7)(t - 1) = 0$ $t = \underline{1, 7/6}$	M1 M1 A1  m1 A1	used allow sign errors any form  method for quad equation Depends on both M's

Q	Solution	Mark	Notes
3(a)(i)	Initial horizontal speed = $15\cos\alpha$ $= 15 \times 0.8$ $= 12 \text{ ms}^{-1}$	B1	
	Time of flight = $9/12$ $= \underline{0.75\text{s}}$	M1 A1	any correct form
3(a)(ii)	Initial vertical speed = $15 \sin\alpha$ $= 15 \times 0.6$ $= 9 \text{ ms}^{-1}$	B1	
	Use of $s = ut + 0.5at^2$ , $u=9(\text{c})$ , $a=(\pm)9.8$ , $t=0.75(\text{c})$ $s = 9 \times 0.75 - 0.5 \times 9.8 \times 0.75^2$ $s = 3.99375 \text{ m}$ Height of B above ground = $\underline{4.99375 \text{ m}}$	M1 A1 A1	si ft s
3(b)	use of $v^2 = u^2 + 2as$ , $u=9$ , $a=(\pm)9.8$ , $s=-1$ $v^2 = 9^2 + 2(-9.8)(-1)$ $v^2 = 100.6$	M1 A1	allow sign errors
	$u_H = 12$	B1	ft candidate's value
	Speed = $\sqrt{12^2 + 100.6}$ Speed = $\underline{15.64 \text{ ms}^{-1}}$	m1 A1	cao

Q	Solution	Mark	Notes
4(a)	Resolve vertically $R \sin \theta = Mg$ $\sin \theta = \frac{3}{5}$ $R = Mg \times \frac{5}{3}$ $R = 5Mg/3$	M1 A1 B1  A1	dim correct    answer given, convincing.
4(b)	N2L towards centre $R \cos \theta = Ma$ $\frac{5Mg}{3} \times \frac{4}{5} = M \times \frac{8g}{3r}$ $CP = r = 2$  $\frac{\text{Height}}{r} = \frac{4}{3}$ $\text{Height} = \frac{8}{3} \text{ m}$	M1 A1  A1  M1  A1	dim correct     use of similar triangles  ft candidate's r if first M1 given.

Q	Solution	Mark	Notes
5(a)	$0 < t < 6$	B1 B1	
5(b)	Distance $t = 6$ to $t = 9 = \int_6^9 2t^2 - 12t \, dt$	M1	use of integration Limits not required
	Distance $= [2t^3/3 - 6t^2]_6^9$ $= 72$	A1	correct integration
	Distance $t = 0$ to $t = 6 = - \int_0^6 2t^2 - 12t \, dt$ Distance $= -[2t^3/3 - 6t^2]_0^6$ $= -[-72]$ $= 72$	A1	or for the other integral
	Required distance $= 72 + 72$ $= \underline{144}$	m1 A1	cao

Q	Solution	Mark	Notes
6(a)	$T = P/v$ $T = \frac{60 \times 1000}{20}$ $T = \underline{3000 \text{ N}}$	M1  A1	used
6(b)	Apply N2L to car and trailer $T - (1500 + 500)g \sin \alpha - (170 + 30) = 2000a$ $3000 - 2000 \times 9.8 \times \frac{1}{14} - 200 = 2000a$ $a = \underline{0.7 \text{ ms}^{-2}}$	M1 A2  A1	dim correct equation All forces present -1 each error  convincing
6(c)	N2L applied to trailer $T - 500g \sin \alpha - 30 = 500a$ $T = 500 \times 9.8 \times \frac{1}{14} + 30 + 500 \times 0.7$ $T = \underline{730 \text{ N}}$ OR N2L applied to car $3000 - 1500g \sin \alpha - 170 - T = 1500 \times 0.7$ $T = 3000 - 1500 \times 9.8 \times \frac{1}{14} - 170 - 1500 \times 0.7$ $T = \underline{730 \text{ N}}$	M1 A2  A1  (M1) (A2)  (A1)	dim correct, all forces -1 each error   dim correct, all forces -1 each error

Q	Solution	Mark	Notes
7(a)	$\text{PE at start} = -2 \times 9.8 \times 0.7$ $= -13.72 \text{ J}$ $\text{PE at end} = -2 \times 9.8 \times (1.2 + x)$ $= -23.52 - 19.6x$ $\text{EE at end} = \frac{1}{2} \times \frac{360}{1.2} x^2$ $\text{EE at end} = 150x^2$ <p>Conservation of energy</p> $150x^2 - 19.6x - 23.52 = -13.72$ $150x^2 - 19.6x - 9.8 = 0$ $x = \underline{0.33}$	M1 A1	mgh used allow 0.7, (1.2+x), (0.5+x), 1.2, 0.5, x.
		M1 A1	use of formula
		M1 A1	equation, all energies correct equation any form
		A1	cao
7(b)	$\text{KE at end} = 0.5 \times 2v^2$ $= v^2$ $\text{PE at end} = -2 \times 9.8 \times 1.2$ $= -23.52$ <p>Conservation of energy</p> $v^2 - 23.52 = -13.72$ $v^2 = 9.8$ $v = \underline{3.13 \text{ ms}^{-1}}$	B1	
		M1 A1	equation, no EE correct equation, any form
		A1	

Q	Solution	Mark	Notes
8(a)	<p>Conservation of energy</p> $0.5mu^2 + mgr\cos\alpha = 0.5mv^2 + mgr\cos\theta$ $0.5 \times 3 \times 5^2 + 3 \times 9.8 \times 4 \times 0.8 =$ $0.5 \times 3 \times v^2 + 3 \times 9.8 \times 4 \times \cos\theta$ $75 + 188.16 = 3v^2 + 235.2\cos\theta$ $v^2 = 87.72 - 78.4\cos\theta$ $v = \sqrt{(87.72 - 78.4\cos\theta)}$	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>equation required</p> <p>KE</p> <p>PE</p> <p>cao</p>
8(b)	<p>N2L towards centre</p> $mg\cos\theta - R = ma$ $R = 3 \times 9.8\cos\theta - \frac{3}{4}(87.72 - 78.4\cos\theta)$ $R = 29.4\cos\theta - 65.79 + 58.8\cos\theta$ $R = \underline{88.2\cos\theta - 65.79}$	<p>M1</p> <p>A1</p> <p>m1</p>	<p>dim correct, all forces</p> <p>substitute, <math>v^2/r</math></p>



### M3

Q	Solution	Mark	Notes
1(a)(i)	Apply N2L to particle $ma = -mg - 3v$ $2 \frac{dv}{dt} = -19.6 - v$	M1 A1	dim correct equation
1(a)(ii)	$\int \frac{2dv}{19.6+v} = - \int dt$ $2\ln 19.6+v  = -t + (C)$ $t = 0, v = 24.5$ $C = 2\ln 44.1 $ $-t = 2\ln\left \frac{19.6+v}{44.1}\right $ $e^{-t/2} = \frac{19.6+v}{44.1}$ $v = 44.1 e^{-t/2} - 19.6$	M1  A1 m1 A1  m1 A1	sep. of variables  correct integration use of initial conditions ft no 2,1/2.  inversion ln to e cao
1(b)	At maximum height, $v = 0$ $t = -2\ln\left \frac{19.6}{44.1}\right $ $= \underline{2 \ln(2.25) = 1.62 \text{ s}}$	M1  A1	si  ft similar expression
1(c)	$\frac{dx}{dt} = 44.1 e^{-t/2} - 19.6$ $x = -88.2 e^{-t/2} - 19.6t (+ C)$ When $t = 0, x = 0$ $C = 88.2$ $x = \underline{88.2 - 88.2 e^{-t/2} - 19.6t}$	M1 A1 m1 A1	$v = \frac{dx}{dt}$ used ft correct integration use of initial conditions ft one slip


Q	Solution	Mark	Notes
2(a)	Amplitude $a = 0.5$	B1	
2(b)	Period $= \frac{2\pi}{\omega} = 2$ $\omega = \pi$ Maximum acceleration $= a\omega^2 = 0.5 \times \pi^2$ Occurs at end points of motion	M1 A1 B1 B1	si ft amplitude $a$ .
2(c)	Let $x = a\cos(\omega t)$ $-0.25 = 0.5\cos(\pi t)$ $\cos(\pi t) = -0.5$ $\pi t = \frac{2\pi}{3}$ $t = \frac{2}{3}$	M1 m1 A1	cao
2(d)	$v^2 = \omega^2(a^2 - x^2), x = 0.3, \omega = \pi$ $v^2 = \pi^2(0.5^2 - 0.3^2)$ $v^2 = \pi^2 \times 0.4^2$ $v = (\pm)0.4\pi$ speed $= 0.4\pi$	M1 A1 A1	ft cao

Q	Solution	Mark	Notes
3(a)(i)	Apply N2L to $P$ $2a = -8x - 10v$ $\frac{d^2x}{dt^2} = -4x - 5\frac{dx}{dt}$	M1 A1	
3(a)(ii)	$\frac{d^2x}{dt^2} + 5\frac{dx}{dt} + 4x = 0$ Auxiliary equation $m^2 + 5m + 4 = 0$ $(m + 4)(m + 1) = 0$ $m = -4, -1$  CF $x = Ae^{-t} + Be^{-4t}$  When $t = 0, x = 2, \frac{dx}{dt} = 3$ $2 = A + B$ $\frac{dx}{dt} = -Ae^{-t} - 4Be^{-4t}$ $3 = -A - 4B$  Adding $5 = -3B$ $B = -\frac{5}{3}$ $A = 2 + \frac{5}{3} = \frac{11}{3}$ $x = \frac{11}{3}e^{-t} - \frac{5}{3}e^{-4t}$	B1 B1 B1 M1 B1 A1 m1 A1	ft values of roots use of initial conditions both equations correct solving simultaneously cao
3(b)	Try $x = at + b$ $\frac{dx}{dt} = a$ $5a + 4(at + b) = 12t - 3$ $4a = 12$ $a = 3$  $5a + 4b = -3$ $15 + 4b = -3$ $4b = -18$ $b = -\frac{9}{2}$  General solution $x = Ae^{-t} + Be^{-4t} + 3t - \frac{9}{2}$	M1 A1 m1 A1	comparing coefficients cao

Q	Solution	Mark	Notes
4	Initial speed of A just before impact = v $v^2 = u^2 + 2as$ , $u=0$ , $a=(\pm)9.8$ , $s=(1.8-0.2)$ $v^2 = 0 + 2 \times 9.8 \times 1.6$ $v = \underline{5.6 \text{ ms}^{-1}}$	M1 A1 A1	cao
	Impulse = Change in momentum Applied to B $J = 3v$	M1 B1	used
	Applied to A $J = 5 \times 5.6 - 5v$	A1	ft c's answer in (a)
	Solving $3v = 28 - 5v$ $8v = 28$ $v = \underline{3.5 \text{ ms}^{-1}}$ $J = \underline{10.5 \text{ N s}}$	m1 A1 A1	cao cao

Q	Solution	Mark	Notes
5(a)	N2L applied to particle		
	$0.25 a = \frac{5}{2x+1}$	M1	
	$v \frac{dv}{dx} = \frac{20}{2x+1}$	M1	$a = v \frac{dv}{dx}$
	$\int v dv = 10 \int \frac{2}{2x+1} dx$	M1	separating variables
	$\frac{1}{2} v^2 = 10 \ln  2x+1  + C$	A1	correct integration ln
	When $x = 0, v = 4$	A1	LHS correct
		m1	use of boundary cond.
			All 3 M's awarded
	$8 = 10 \ln(1) + C$		
	$C = 8$		
	$v^2 = 20 \ln  2x+1  + 16$		
	$\ln  2x+1  = \frac{1}{20} (v^2 - 16)$		
	$2x+1 = e^{0.05(v^2-16)}$	m1	inversion, 3 M's awarded
	$x = 0.5(e^{0.05(v^2-16)} - 1)$	A1	cao any equivalent exp.
5(b)	$v = 6$		
	$x = 0.5(e^{0.05(36-16)} - 1)$	M1	exp. with $v^2$ needed
	$x = 0.5(e - 1)$		
	$x = \underline{0.86 \text{ m}}$	A1	cao
5(c)	$a = 5$		
	$\frac{20}{2x+1} = 5$	M1	
	$20 = 10x + 5$		
	$x = 1.5$	A1	
	$v^2 = 20 \ln(3+1) + 16$	m1	substitution in expression with $v^2$ .
	$= 20 \ln 4 + 16$		
	$v = \underline{6.61 \text{ ms}^{-1}}$	A1	cao

Q	Solution	Mark	Notes
---	----------	------	-------



6

6(a)	Moments about A	M1	equation, no extra forces No missing forces
	$6g \times 2 + 3g \times 3 = T \times 4 \sin \theta$	A2	-1 each error
	$4 \times \frac{3}{5} T = 21g$		
	$T = \frac{35}{4} g = 85.75 \text{ N}$	A1	cao
6(b)	Resolve vertically	M1	equation, all forces, no extra force
	$T \sin \theta + Y = 9g$	A1	
	$Y = 9g - \frac{35}{4} g \times \frac{3}{5}$		
	$Y = \frac{15}{4} g = 36.75 \text{ N}$	A1	cao
	Resolve horizontally		
	$T \cos \theta = X$	M1	equation, all forces, No extra force
	$X = \frac{35}{4} g \times \frac{4}{5}$		
	$X = 7g = 68.6 \text{ N}$	A1	cao
6(b)(i)	Magnitude of reaction at wall		
	$= \sqrt{68.6^2 + 36.75^2}$	M1	
	$= 77.82 \text{ N}$	A1	ft X and Y
6(b)(ii)	$\mu = \frac{Y}{X}$	M1	used
	$\mu = \frac{15}{4 \times 7} = \frac{15}{28}$	A1	ft X and Y if answer < 1.

Ques	Solution	Mark	Notes
1(a)	$P(A \cup B) = P(A) + P(B)$ $P(B) = 0.4 - 0.25 = 0.15$	M1 A1	Award M1 for using formula
(b)	$P(A \cup B) = P(A) + P(B) - P(A)P(B)$ $0.4 = 0.25 + P(B) - 0.25P(B)$ $P(B) = 0.15/0.75 = 0.2$	M1 A1 A1	Award M1 for using formula
2(a)	$P(1 \text{ of each}) =$ $\frac{5}{10} \times \frac{3}{9} \times \frac{2}{8} \times 6 \text{ or } \binom{5}{1} \times \binom{3}{1} \times \binom{2}{1} \div \binom{10}{3}$ $= \frac{1}{4}$	M1A1 A1	M1A0A0 if 6 omitted Special case : if they use an incorrect total, eg 9 or 11, FT their incorrect total but subtract 2 marks at the end
(b)	$P(3 \text{ war}) = \frac{5}{10} \times \frac{4}{9} \times \frac{3}{8} \text{ or } \binom{5}{3} \div \binom{10}{3}$ $= \frac{1}{12}$	M1 A1	
(c)	$P(3 \text{ cowboy}) = \frac{3}{10} \times \frac{2}{9} \times \frac{1}{8} \text{ or } \binom{3}{3} \div \binom{10}{3}$ $= \frac{1}{120}$ $P(3 \text{ the same}) = \frac{1}{12} + \frac{1}{120} = \frac{11}{120}$	B1 M1A1	FT previous values
3	$E(X) = 20$ $\text{Var}(X) = 4 \text{ (SD} = 2)$ $E(Y) = 20a + b = 65$ $\text{Var}(Y) = 4a^2 = 36$ $a = 3$ $b = 5$	B1 B1 B1 B1 B1 B1	Accept $\text{SD}(Y) = 2a = 6$ Must be justified by solving the two equations
4(a)(i)	$B(20, 0.25)$	B1	B must be mentioned and the parameters $n$ and $p$ must be seen or implied somewhere in the question
(ii)	$P(3 \leq X \leq 9) = 0.9087 - 0.0139 \text{ or } 0.9861 - 0.0913$ $= 0.8948$	B1B1 B1	FT an incorrect $p$ except for the last three marks M0 if no working seen
(iii)	$P(X = 6) = \binom{20}{6} \times 0.25^6 \times 0.75^{14}$ $= 0.169$	M1 A1	
(b)(i)	Let $Y$ denote the number of throws giving '8' Then $Y$ is $B(160, 0.0625) \approx \text{Poi}(10)$ . $P(Y = 12) = e^{-10} \times \frac{10^{12}}{12!}$ $= 0.0948$	B1 M1 A1	M0 if no working seen Accept the use of tables Correct values only (no FT)
(ii)	$P(6 \leq Y \leq 14) = 0.9165 - 0.0671 \text{ or } 0.9329 - 0.0835$ $= 0.8494 \text{ cao}$	B1B1 B1	

<b>5(a)</b>	$P(1) = \frac{1}{3} \times \frac{1}{4} + \frac{1}{3} \times \frac{1}{3} + \frac{1}{3} \times \frac{1}{2}$ $= \frac{13}{36} \quad (0.361)$	<b>M1A1</b> <b>A1</b>	M1 Use of Law of Total Prob (Accept tree diagram)
<b>(b)</b>	$P(A 1) = \frac{1/12}{13/36}$ $= \frac{3}{13} \quad \text{cao} \quad (0.231)$	<b>B1B1</b> <b>B1</b>	FT denominator from (a) B1 num, B1 denom
<b>6(a)</b>	The sequence is MMMH si Prob = $0.3 \times 0.3 \times 0.3 \times 0.7 = 0.0189$	<b>B1</b> <b>M1A1</b>	Award B1 for 0.147
<b>(b)</b>	The sequence is MHH or HMH si Prob = $0.3 \times 0.7 \times 0.7 + 0.7 \times 0.3 \times 0.7 = 0.294$	<b>B1</b> <b>M1A1</b>	
<b>7(a)</b>	$\sum p_x = k \left( 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} \right) = 1$ $k \left( \frac{8+4+2+1}{8} \right) = 1 \rightarrow k = \frac{8}{15}$	<b>M1</b> <b>A1</b>	Convincing
<b>(b)</b>	$E(X) = \frac{8}{15} \times 1 + \frac{4}{15} \times 2 + \frac{2}{15} \times 4 + \frac{1}{15} \times 8$ $= \frac{32}{15} \quad (2.13)$ $E(X^2) = \frac{8}{15} \times 1 + \frac{4}{15} \times 4 + \frac{2}{15} \times 16 + \frac{1}{15} \times 64 \quad (8)$	<b>M1</b> <b>A1</b> <b>M1A1</b>	
<b>(c)(i)</b>	$\text{Var}(X) = 8 - \left( \frac{32}{15} \right)^2 = 3.45 \quad (776/225)$ <p>The possibilities are (1,1); (2,2); (4,4); (8,8) si</p> $P(X_1 = X_2) = \left( \frac{8}{15} \right)^2 + \left( \frac{4}{15} \right)^2 + \left( \frac{2}{15} \right)^2 + \left( \frac{1}{15} \right)^2$ $= \frac{17}{45} \quad (0.378)$	<b>A1</b> <b>B1</b> <b>M1</b>	Accept 3.46
<b>(ii)</b>	<p>It follows that <math>P(X_1 \neq X_2) = \frac{28}{45}</math></p> <p>And therefore by symmetry <math>P(X_1 &gt; X_2) = \frac{14}{45}</math></p>	<b>A1</b> <b>M1</b> <b>A1</b>	
			FT their answer from (c)(i) Do not accept any other method.



<p><b>8(a)</b></p> <p><b>(b)</b></p>	<p>Let <math>X</math> denote the number of calls between 9am and 10 am so that <math>X</math> is <math>Po(5)</math></p> $P(X = 7) = \frac{e^{-5} \times 5^7}{7!}$ $= 0.104$ <p>We require</p> $P(\text{calls betw 9 and 10}=7   \text{calls betw 9 and 11}=10)$ $= \frac{P(c \text{ b } 9 \text{ and } 10 = 7 \text{ AND } c \text{ b } 9 \text{ and } 11 = 10)}{P(\text{calls between 9 and 11} = 10)}$ $= \frac{P(c \text{ b } 9 \text{ and } 10 = 7) \times P(c \text{ b } 10 \text{ and } 11 = 3)}{P(\text{calls between 9 and 11} = 10)}$ $= \frac{e^{-5} \times 5^7}{7!} \times \frac{e^{-5} \times 5^3}{3!} \div \frac{e^{-10} \times 10^{10}}{10!} \quad (\text{denom} = 0.125)$ $= 0.117$	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1A1</b></p> <p><b>A1</b></p>	<p>M0 no working</p> <p>A1 numerator, A1 denominator The denominator A1 can be awarded if the M1 is awarded</p>
<p><b>9(a)</b></p> <p><b>(b)</b></p> <p><b>(c)(i)</b></p> <p><b>(ii)</b></p>	$\int_0^2 k \left( 1 - \frac{x^2}{4} \right) dx = 1$ $k \left[ x - \frac{x^3}{12} \right]_0^2 = 1$ $k \left( 2 - \frac{8}{12} \right) = 1$ $k = \frac{3}{4}$ $E(X) = \int_0^2 x \left( \frac{3}{4} - \frac{3x^2}{16} \right) dx$ $= \left[ \frac{3x^2}{8} - \frac{3x^4}{64} \right]_0^2$ $= 0.75$ $F(x) = \int_0^x \left( \frac{3}{4} - \frac{3t^2}{16} \right) dt$ $= \left[ \frac{3t}{4} - \frac{t^3}{16} \right]_0^x$ $= \frac{3x}{4} - \frac{x^3}{16}$ $P(0.5 \leq X \leq 1.5) = F(1.5) - F(0.5)$ $= 0.547$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>M1 for <math>\int f(x)dx</math>, limits not required until next line</p> <p>M1 for the integral of <math>xf(x)</math>, A1 for completely correct although limits may be left until 2<sup>nd</sup> line.</p> <p>M1 for <math>\int f(x)dx</math></p> <p>A1 for performing the integration</p> <p>A1 for dealing with the limits</p> <p>FT their <math>F(x)</math></p>

Ques	Solution	Mark	Notes
1(a)(i)	$z = \frac{10.5 - 10}{2} = 0.25$ $P(X \leq 10.5) = 0.5987$	M1A1 A1	M0 for $2^2$ or $\sqrt{2}$ M1A0 for $-0.25$ if final answer incorrect M0 no working
(ii)	$x = \frac{x - \mu}{\sigma} = 1.282$ $= 12.564$	M1 A1	M1 for 2.326, 1.645, 2.576 Accept 12.6
(b)(i)	$E(X + 2Y) = 34$ $\text{Var}(X + 2Y) = \text{Var}(X) + 4\text{Var}(Y)$ $= 40$ <p>We require <math>P(X + 2Y &lt; 36)</math></p> $z = \frac{36 - 34}{\sqrt{40}} = 0.32$	B1 B1 M1A1	FT their mean and variance M0 no working
(ii)	$\text{Prob} = 0.6255$ <p>Consider <math>U = X_1 + X_2 + X_3 - Y_1 - Y_2</math></p> $E(U) = 3 \times 10 - 2 \times 12 = 6$ $\text{Var}(U) = 3 \times 4 + 2 \times 9 = 30$ <p>We require <math>P(U &lt; 0)</math></p> $z = \frac{0 - 6}{\sqrt{30}} = -1.10$ $\text{Prob} = 0.136$	A1 B1 M1A1 m1A1 A1	Do not FT their mean and variance
2(a)	$\bar{x} = \frac{9980}{50} (= 199.6)$ $\text{SE of } \bar{X} = \frac{4}{\sqrt{50}} (= 0.5656\dots)$ <p>95% conf limits are  <math>199.6 \pm 1.96 \times 0.5656\dots</math>  giving [198.5, 200.7] cao</p>	B1 B1 M1A1 A1	M1 correct form, A1 correct z. M0 no working
(b)	<p>Width of 95% CI = <math>3.92 \times \frac{4}{\sqrt{n}}</math> si</p> <p>We require</p> $3.92 \times \frac{4}{\sqrt{n}} < 1$ $n > 245.86\dots$ <p>Minimum <math>n = 246</math></p>	B1 M1 A1 A1	FT their z from (a)  Award M1A0A0 for 1.96 instead of 3.92 FT from line above if $n > 50$

<b>3(a)</b>	$H_0 : \mu_B = \mu_G; H_1 : \mu_B \neq \mu_G$	<b>B1</b>	
<b>(b)</b>	$\bar{x}_B = \frac{482}{8} = 60.25; \bar{x}_G = \frac{430}{8} = 53.75$ SE of diff of means = $\sqrt{\frac{7.5^2}{8} + \frac{7.5^2}{8}} \quad (3.75)$ Test statistic (z) = $\frac{60.25 - 53.75}{3.75}$ = 1.73 Prob from tables = 0.0418 p-value = 0.0836 Insufficient evidence to conclude that there is a difference in performance between boys and girls.	<b>B1B1</b>  <b>M1A1</b>  <b>m1A1</b>  <b>A1</b> <b>A1</b> <b>B1</b>  <b>B1</b>	     FT their z if M marks gained FT on line above  FT their p-value
<b>4(a)</b>	$H_0 : p = 0.4; H_1 : p > 0.4$	<b>B1</b>	
<b>(b)</b>	Let X = No. supporting politician so that X is B(50,0.4) (under $H_0$ ) si p-value = $P(X \geq 25   X \text{ is B}(50,0.4))$ = 0.0978 Insufficient evidence to conclude that the support is greater than 40%.	<b>B1</b> <b>M1</b> <b>A1</b>  <b>B1</b>	  M0 for $P(X = 25)$ or $P(X > 25)$ M0 normal or Poisson approx  FT on p-value
<b>(c)</b>	X is now B(400,0.4) (under $H_0$ ) $\approx N(160,96)$ p-value = $P(X \geq 181   X \text{ is } N(160,96))$ $z = \frac{180.5 - 160}{\sqrt{96}}$ = 2.09 p-value = 0.0183 Strong evidence to conclude that the support is greater than 40%.	<b>B1</b> <b>M1</b>  <b>m1A1</b>  <b>A1</b> <b>A1</b>  <b>B1</b>	   Award m1A0A1A1 for incorrect or no continuity correction 181.5 $\rightarrow z = 2.19 \rightarrow p = 0.01426$ 181 $\rightarrow z = 2.14 \rightarrow p = 0.01618$  FT on p-value
<b>5(a)</b>	$H_0 : \mu = 1.2 : H_1 : \mu < 1.2$	<b>B1</b>	Must be $\mu$
<b>(b)(i)</b>	Let X = number of accidents in 60 days Then X is Poi(72) (under $H_0$ ) $\approx N(72,72)$ si  Sig level = $P(X \leq 58   H_0)$ $z = \frac{58.5 - 72}{\sqrt{72}}$ = -1.59 Sig level = 0.0559	<b>B1</b>  <b>M1</b> <b>m1A1</b>  <b>A1</b> <b>A1</b> <b>B1</b> <b>M1</b>	  Award m1A0A1A1 for incorrect or no continuity correction 57.5 $\rightarrow z = -1.71 \rightarrow p = 0.0436$ 58 $\rightarrow z = -1.65 \rightarrow p = 0.0495$
<b>(ii)</b>	X is now Poi(48) which is approx N(48,48) si P(wrong conclusion) = $P(X \geq 59   \mu = 0.8)$ $z = \frac{58.5 - 48}{\sqrt{48}}$ = 1.52 P(wrong conclusion) = 0.0643	<b>B1</b>  <b>m1A1</b> <b>A1</b> <b>A1</b>	  Award m1A0A1A1 for incorrect or no continuity correction 59.5 $\rightarrow z = 1.66 \rightarrow p = 0.0485$ 59 $\rightarrow z = 1.59 \rightarrow p = 0.0559$

<b>6(a)(i)</b>	$E(C) = 2\pi E(R)$ $= 2\pi \times 7 = 14\pi \quad (43.98)$ $\text{Var}(C) = 4\pi^2 \text{Var}(R)$ $= \frac{4\pi^2}{3} \quad (13.16)$	<b>M1</b> <b>A1</b> <b>M1</b>	Accept the use of integration, M1 for a correct integral and A1 for the correct answer
<b>(ii)</b>	$P(C \leq 45) = P(R \leq 45/2\pi)$ $= \frac{(45/2\pi - 6)}{8 - 6}$ $= 0.581$	<b>M1</b> <b>A1</b>  <b>A1</b>	
<b>(b)(i)</b>	$A = \pi R^2$ $P(A \geq 150) = P\left(R \geq \sqrt{150/\pi}\right)$ $= \frac{8 - \sqrt{150/\pi}}{8 - 6}$	<b>M1A1</b>  <b>A1</b>	
<b>(ii)</b>	<p>EITHER</p> $E(A) = \int_6^8 \pi r^2 \times \frac{1}{2} dr$ $= \frac{\pi}{6} \left[ r^3 \right]_6^8$ $= \frac{148\pi}{3} \quad (155)$ <p>OR</p> $E(A) = \pi E(R^2) = \pi (\text{var}(R) + (E(R))^2)$ $= \pi \left( \frac{1}{3} + 7^2 \right)$ $= \frac{148\pi}{3} \quad (155)$	 <b>A1</b>  <b>M1</b>  <b>A1</b>  <b>A1</b>  <b>M1</b>  <b>A1</b>  <b>A1</b>	

**S3**

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>								
<b>1</b>	$\hat{p} = 0.29 \text{ si}$ $\text{ESE} = \sqrt{\frac{0.29 \times 0.71}{300}} (= 0.02619..) \text{ si}$ 95% confidence limits are $0.29 \pm 1.96 \times 0.02619..$ giving [0.24, 0.34]	<b>B1</b>  <b>M1A1</b>  <b>m1A1</b> <b>A1</b>	m1 correct form, A1 1.96								
<b>2</b>	<p>The possibilities are  <u>3 red, 1 blue for which <math> X - Y  = 2</math></u>            Therefore,</p> $P(X - Y = 2) = \frac{3}{10} \times \frac{2}{9} \times \frac{1}{8} \times \frac{7}{7} \times 4 \text{ OR } \frac{\binom{3}{3} \times \binom{7}{1}}{\binom{10}{4}}$ $= \frac{1}{30}$ <p><u>2 red, 2 blue for which <math> X - Y  = 0</math></u></p> $P(X - Y = 0) = \frac{3}{10} \times \frac{2}{9} \times \frac{7}{8} \times \frac{6}{7} \times 6 \text{ OR } \frac{\binom{3}{2} \times \binom{7}{2}}{\binom{10}{4}}$ $= \frac{3}{10}$ <p><u>1 red, 3 blue for which <math> X - Y  = 2</math></u></p> $P(X - Y = -2) = \frac{3}{10} \times \frac{7}{9} \times \frac{6}{8} \times \frac{5}{7} \times 4 \text{ OR } \frac{\binom{3}{1} \times \binom{7}{3}}{\binom{10}{4}}$ $= \frac{1}{2}$ <p><u>0 red, 4 blue for which <math> X - Y  = 4</math></u></p> $P(X - Y = -4) = \frac{7}{10} \times \frac{6}{9} \times \frac{5}{8} \times \frac{4}{7} \text{ OR } \frac{\binom{7}{4}}{\binom{10}{4}} = \frac{1}{6}$ <p>The distribution of <math> X - Y </math> is therefore</p> <table border="1"> <tr> <td><math> X - Y </math></td><td>0</td><td>2</td><td>4</td></tr> <tr> <td>Prob</td><td>3/10</td><td>8/15</td><td>1/6</td></tr> </table>	$ X - Y $	0	2	4	Prob	3/10	8/15	1/6	<b>M1A1</b>   <b>A1</b>   <b>B1</b>   <b>B1</b>   <b>B1</b>   <b>M1A1</b>	FT if found as $1 - \Sigma \text{probs}$       FT their probabilities
$ X - Y $	0	2	4								
Prob	3/10	8/15	1/6								

<p><b>3(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	<p>UE of <math>\mu = 34.3</math>  <math>\Sigma x^2 = 10609.43</math>            UE of <math>\sigma^2 = \frac{10609.43}{8} - \frac{9 \times 34.3^2}{8}</math>  <math>= 2.6275</math>            DF = 8 si            t-value = 1.86            90% confidence limits are  <math>34.3 \pm 1.86 \sqrt{\frac{2.6275}{9}}</math>            giving [33.3, 35.3] cao</p> <p>EITHER            Width of interval = <math>2t \sqrt{\frac{2.6275}{9}} = 3.2</math>            So <math>t = 2.96</math>            For a 99% confidence interval, <math>t = 3.355</math>            Since <math>2.96 &lt; 3.355</math>, the confidence level is less than 99%            OR            For 99% confidence interval, <math>t = 3.355</math>            99% confidence limits are  <math>34.3 \pm 3.355 \sqrt{\frac{2.6275}{9}}</math>            giving [32.5, 36.1]            The given confidence interval is narrower than this therefore its confidence level is less than 99%</p>	<p><b>B1</b> <b>B1</b></p> <p><b>M1</b> <b>A1</b></p> <p><b>B1</b> <b>B1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p><b>M1</b> <b>A1</b> <b>B1</b> <b>A1</b></p> <p><b>B1</b></p> <p><b>M1</b> <b>A1</b></p> <p><b>A1</b></p>	<p>No working need be seen</p> <p>M0 division by 9 Answer only no marks</p> <p>Answer only no marks</p>
<p><b>4(a)</b></p> <p><b>(b)</b></p>	<p>The 5% critical value = <math>2000 + 1.645 \times \sqrt{\frac{2554}{120}}</math>  <math>= 2007.6</math>            The 10% critical value = <math>2000 + 1.282 \times \sqrt{\frac{2554}{120}}</math>  <math>= 2005.9</math>            The required range is therefore            (2005.9, 2007.6)            No because of the Central Limit Theorem            AND THEN EITHER            which ensures the normality of the sample mean            OR            which can be used because the sample is large</p>	<p><b>M1</b> <b>A1</b></p> <p><b>M1</b> <b>A1</b></p> <p><b>A1</b> <b>B1</b></p> <p><b>B1</b></p>	<p>M1A0 for –</p> <p>M1A0 for –</p>



<p><b>7(a)</b></p>	$E(\hat{p}) = \frac{E(X)}{n} = \frac{np}{n} = p$ <p>Therefore unbiased.</p> $SE(\hat{p}) = \sqrt{\frac{\text{Var}(X)}{n^2}} = \sqrt{\frac{np(1-p)}{n^2}} = \sqrt{\frac{p(1-p)}{n}}$	<p><b>M1</b> <b>A1</b></p>	<p>This line need not be seen</p>
<p><b>(b)(i)</b></p>	$E(\hat{p}^2) = \frac{E(X^2)}{n^2}$ $= \frac{\text{Var}(X) + [E(X)]^2}{n^2}$ $= \frac{np(1-p) + n^2 p^2}{n^2}$ $= \left( p^2 + \frac{p(1-p)}{n} \right)$	<p><b>M1</b> <b>m1</b> <b>A1</b></p>	<p>Accept <math>q</math> for <math>1-p</math></p>
<p><b>(ii)</b></p>	$\neq p^2 \text{ therefore not unbiased}$ $E[X(X-1)] = E(X^2) - E(X)$ $= np(1-p) + n^2 p^2 - np$ $= n(n-1)p^2$ <p>It follows that</p> $\frac{X(X-1)}{n(n-1)}$	<p><b>A1</b> <b>M1</b> <b>A1</b> <b>A1</b> <b>A1</b></p>	<p>This line need not be seen</p>
<p><b>(c)(i)</b></p>	<p>is an unbiased estimator for <math>p^2</math>.</p> <p>EITHER</p> <p>By reversing the interpretation of success and failure, it follows that</p> $\frac{(n-X)(n-X-1)}{n(n-1)}$ <p>is an unbiased estimator for <math>q^2</math>.</p> <p>OR</p> $q^2 = (1-p)^2 = 1 - 2p + p^2$ <p>Therefore an unbiased estimator for <math>q^2</math> is</p>	<p><b>M1</b> <b>A1</b> <b>M1</b></p>	
<p><b>(ii)</b></p>	$1 - \frac{2X}{n} + \frac{X(X-1)}{n(n-1)}$ <p>Since <math>pq = p(1-p) = p - p^2</math></p> <p>It follows that an unbiased estimator for <math>pq</math></p> $= \frac{X}{n} - \frac{X(X-1)}{n(n-1)}$ $= \frac{X(n-X)}{n(n-1)}$	<p><b>A1</b> <b>M1</b> <b>A1</b> <b>A1</b></p>	<p>This expression need not be simplified</p>





WJEC  
245 Western Avenue  
Cardiff CF5 2YX  
Tel No 029 2026 5000  
Fax 029 2057 5994  
E-mail: [exams@wjec.co.uk](mailto:exams@wjec.co.uk)  
website: [www.wjec.co.uk](http://www.wjec.co.uk)