


# M1

| Q | Solution | Mark | Notes |
|---|----------|------|-------|
|---|----------|------|-------|

1.



N2L applied to man

M1  $R$  and  $Mg$  opposing.  
dim correct

$$R - Mg = Ma$$

A1

$$680 = M(9.8 + 0.2)$$

$$M = \underline{68}$$

A1 cao

N2L applied to Lift and Man

M1  $T$  and weight opposing.  
dim correct.

$$T - 1868g = 1868a$$

A1 ft  $M$

$$T = \underline{18680 \text{ (N)}}$$

A1 ft  $M$


| Q  | Solution                                                                   | Mark | Notes                                    |
|----|----------------------------------------------------------------------------|------|------------------------------------------|
| 2. | Apply N2L to $B$                                                           | M1   | dim correct, all forces.                 |
|    | $5g - T = 0$                                                               | A1   | allow $5a$ RHS<br>$5g$ and $T$ opposing. |
|    | Resolve perpendicular to plane for $A$                                     | M1   | allow sin                                |
|    | $R = 4g \cos \alpha$                                                       | A1   |                                          |
|    | Apply N2L to $A$                                                           | M1   | Friction opposes motion.                 |
|    | $T - 4g \sin \alpha - F = 0$                                               | A1   | Allow $4a$ RHS and/or cos                |
|    | At limiting equilibrium $F = \mu R$                                        | M1   | used                                     |
|    | $\mu = \frac{F}{R} = \frac{45g}{48g} = \frac{15}{16}$                      | A1   | convincing                               |
|    | $T = 5g = 49$                                                              |      |                                          |
|    | $F = T - 4g \sin \alpha = \frac{45g}{13} = \frac{441}{13} = 33.9231$       |      |                                          |
|    | $R = 4g \times \frac{12}{13} = \frac{48g}{13} = \frac{2352}{65} = 36.1846$ |      |                                          |

| Q    | Solution                                                    | Mark | Notes                             |
|------|-------------------------------------------------------------|------|-----------------------------------|
| 3(a) | Conservation of momentum                                    | M1   | attempted, equation, dim correct. |
|      | $3 \times 8 + 5 \times 2 = 3v_A + 5v_B$                     | A1   |                                   |
|      | $3v_A + 5v_B = 34$                                          |      |                                   |
|      | Restitution                                                 | M1   |                                   |
|      | $v_B - v_A = -\frac{1}{3}(2 - 8)$                           | A1   |                                   |
|      | $v_B - v_A = 2$                                             |      |                                   |
|      | $3v_A + 5v_B = 34$                                          |      |                                   |
|      | $-3v_A + 3v_B = 6$                                          |      |                                   |
|      | Adding                                                      | m1   | dep on both M's                   |
|      | $8v_B = 40$                                                 |      |                                   |
|      | $v_B = \underline{5 \text{ (ms}^{-1}\text{)}}$              | A1   | cao                               |
|      | $v_A = \underline{3 \text{ (ms}^{-1}\text{)}}$              | A1   | cao                               |
| 3(b) | Impulse = change of momentum                                | M1   | used                              |
|      | $I = 5 \times 5 - 5 \times 2 = \underline{15 \text{ (Ns)}}$ | A1   | ft $v_A$ or $v_B$                 |

| Q | Solution                                                                                                               | Mark                                       | Notes |
|---|------------------------------------------------------------------------------------------------------------------------|--------------------------------------------|-------|
| 4 | <p>Moments about <math>x</math>-axis</p> $= 5 \times (-1) + 2 \times (3) + 3 \times 5 + 6 \times 0$ $16y = 16$ $y = 1$ | <p>B1</p> <p>M1    si</p> <p>A1    cao</p> |       |
|   | <p>Moments about <math>y</math>-axis</p> $= 5 \times 4 + 2 \times 2 + 3 \times (-2) + 6 \times (-3)$ $16x = 0$ $x = 0$ | <p>B1</p> <p>M1    si</p> <p>A1    cao</p> |       |

| Q | Solution | Mark | Notes |
|---|----------|------|-------|
|---|----------|------|-------|

5(a)



Moments about A

$$2.8R_B = 80g \times 0.9 + 15g \times 1.4$$

$$R_B = \underline{325.5 \text{ (N)}}$$

Vertical forces in equilibrium

$$R_A + R_B = 80g + 15g$$

$$R_A = \underline{605.5 \text{ (N)}}$$

M1 3 terms, dim correct  
Equation required  
A1 correct equation  
B1 any correct moment


A1 cao

M1 all forces, no extra

A1

A1 cao

5(b)



Resolve vertically

$$1.5R + R = 95g$$

$$R = 38g$$

Moments about A

$$2.8 \times R = 80g \times 0.9 + 15g \times x$$

$$x = \frac{172}{75} = \underline{2.3 \text{ (m)}}$$

M1

A1

M1 3 terms, dim correct

A1 oe

A1 cao

| Q    | Solution                                                                                                                       | Mark                                                                                        | Notes |
|------|--------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|-------|
| 6(a) |                                                                                                                                | <p>B1 labels, units and shape<br/> B1 (0, 0) to (10, 20)<br/> B1 (10, 20) to (10+T, 20)</p> |       |
| 6(b) | $v = u + at, v=20, u=0, t=10$<br>$20 = 0 + 10a$<br>$a = \underline{2 \text{ (ms}^{-2}\text{)}}$                                | <p>M1<br/> A1</p>                                                                           |       |
| 6(c) | <p>Total distance = area under graph</p> $D = 0.5 \times 10 \times 20 + 20T$<br>$D = 100 + 20T \text{ (m)}$                    | <p>M1 attempted<br/> B1 one correct area<br/> A1 cao</p>                                    |       |
| 6(d) | $s = ut + 0.5at^2, u=0, t=5+T, a=2$<br>$s = 0.5 \times 2 \times (5+T)^2$<br>$D = 25 + 10T + T^2$                               | <p>M1<br/> A1</p>                                                                           |       |
|      | $25 + 10T + T^2 = 100 + 20T$<br>$T^2 - 10T - 75 = 0$<br>$(T + 5)(T - 15) = 0$<br>$T = 15$<br>$D = \underline{400 \text{ (m)}}$ | <p>M1 Ft exp for D in (d) and (c)<br/> A1 cao<br/> A1 cao</p>                               |       |

| Q      | Solution                                                            | Mark     | Notes                       |
|--------|---------------------------------------------------------------------|----------|-----------------------------|
| 7      | Resolve in 80 N direction<br>$80 = P\cos 60^\circ + Q\cos 45^\circ$ | M1<br>A1 | Equation required           |
|        | Resolve in 25 N direction<br>$25 = P\sin 60^\circ - Q\sin 45^\circ$ | M1<br>A1 | Equation required           |
|        | $160 = P + Q\sqrt{2}$<br>$50 = P\sqrt{3} - Q\sqrt{2}$               |          |                             |
| Adding |                                                                     | m1       | dep on both M's             |
|        | $(1 + \sqrt{3})P = 210$                                             |          |                             |
|        | $P = \underline{76.9}$                                              | A1       | cao                         |
|        | $Q = \underline{58.8}$                                              | A1       | cao                         |
|        |                                                                     |          | penalise once if not 1 d.p. |

| Q    | Solution                                                                                                                                                                                                                    | Mark                           | Notes                     |
|------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|---------------------------|
| 8(a) | Use of $v^2=u^2+2as$ with $u=(\pm)2.1, a=(\pm)9.8,$<br>$s=(\pm)4.$<br>$v^2 = 2.1^2 + 2 \times 9.8 \times 4$<br>$v = 9.1$<br><br>speed of rebound $= 9.1 \times \frac{4}{7}$<br>$= \underline{5.2 \text{ (ms}^{-1}\text{)}}$ | M1<br>A1<br>A1<br><br>m1<br>A1 | allow -<br><br>convincing |
| 8(b) | We require smallest $n$ st $\left(\frac{4}{7}\right)^n \times 9.1 < 1$<br><br>4 bounces                                                                                                                                     | M1<br><br>A1                   | oe, si trial & error      |



| Q | Solution                                                    |            |     |     | Mark | Notes                                                         |
|---|-------------------------------------------------------------|------------|-----|-----|------|---------------------------------------------------------------|
| 9 | <i>BCD</i>                                                  | 45         | 19  | (5) | B1   | for 19                                                        |
|   | <i>ABDE</i>                                                 | 160        | 8   | (5) |      |                                                               |
|   | Circle                                                      | $9\pi$     | 7   | (5) | B1   | both 8 and 7 required                                         |
|   | Lamina                                                      | $205-9\pi$ | $x$ | (y) | B1   | expressions for areas, oe                                     |
|   | Moments about <i>AE</i>                                     |            |     |     | M1   |                                                               |
|   | $(205-9\pi)x + 9\pi \times 7 = 160 \times 8 + 45 \times 19$ |            |     |     | A1   | signs correct. Ft table if at least one B1 for c of m gained. |
|   | $x = \underline{10.96}$                                     |            |     |     | A1   | cao                                                           |
|   |                                                             |            |     |     |      |                                                               |
|   | $y = \underline{5}$                                         |            |     |     | B1   |                                                               |
|   |                                                             |            |     |     |      |                                                               |